

TOPIC

Temi di Psicologia dell'Ordine degli Psicologi
della Campania

NEUROPSICOLOGIA E NEUROSCIENZE

· Conseguenze emotive a seguito di lesioni cerebrali acquisite: una revisione narrativa della letteratura

- Aurora Leontina Lombardi, Michele Bruno, Giuseppina De Rosa, Roberta Ranieri, Luca Santella, Roberta Beatrice Sarnataro, Valentina Sartorio, Gabriella Santangelo, Alfonsina D'Iorio

PSICOLOGIA DELLO SVILUPPO

· Gli effetti di un intervento di psicologia positiva sul benessere psicologico degli adolescenti: uno studio preliminare

- Mariachiara Gaita, Mariamichela Aquino, Alfredo di Dio, Maria Cropano, Sabatino Galietta, Antonella Fiorentino, Simona Raimo, Florindo d'Onofrio

Emotional sequelae following acquired brain injury: a narrative review.

Conseguenze emotive a seguito di lesioni cerebrali acquisite: una revisione narrativa della letteratura

Aurora Leontina Lombardi¹, Michele Bruno¹, Giuseppina De Rosa¹, Roberta Ranieri¹, Luca Santella¹,
Roberta Beatrice Sarnataro¹, Valentina Sartorio¹, Gabriella Santangelo¹, Alfonsina D'Iorio¹

1. Dipartimento di Psicologia, Università degli Studi della Campania "Luigi Vanvitelli", Caserta, Italia

Abstract

Le lesioni cerebrali acquisite (LCA), che comprendono traumi cranici (TBI) e ictus, comportano una rosa di conseguenze emotive che possono incidere profondamente sul funzionamento sociale e sulla qualità della vita dei pazienti. Questa revisione narrativa vuole riassumere le conseguenze emotive più di frequente associate alle LCA, tra cui ansia, depressione, apatia, anedonia, disinibizione, aggressività e deficit nella teoria della mente. La depressione è caratterizzata da pervasivi sentimenti negativi soggettivi e da una sofferenza emotiva, spesso accompagnata da una riduzione della motivazione e da una tendenza al ritiro dalle attività quotidiane. Al contrario, l'ansia può comportare una maggiore sensibilità agli stimoli interni ed esterni, che porta ad una persistente sensazione di tensione e attivazione fisiologica, che può anch'essa influire sulla motivazione. Sebbene apatia e anedonia possano sovrapporsi in parte con cambiamenti della motivazione, sono concetti distinti: l'apatia è definita da indifferenza emotiva e distacco affettivo, mentre l'anedonia si riferisce ad una capacità ridotta di provare piacere o interesse per attività precedentemente gratificanti. La disregolazione emotiva, come l'aumento dell'irritabilità, l'aggressività e i comportamenti impulsivi, può contribuire all'isolamento sociale. La disinibizione può manifestarsi attraverso comportamenti sociali inappropriati, complicando la reintegrazione sociale dei pazienti. Infine, i deficit nella teoria della mente, ovvero nella capacità di comprendere le emozioni e le prospettive altrui, possono ostacolare le interazioni e le relazioni sociali. Questa revisione narrativa della letteratura vuole evidenziare la natura pervasiva e multifattoriale delle conseguenze emotive risultanti dalle LCA. Tali disturbi sono spesso persistenti e complessi, si vuole quindi sottolineare la necessità di una maggiore attenzione e dell'implementazione di programmi di riabilitazione personalizzati per migliorare la qualità della vita dei pazienti.

Parole chiave

Lesione Cerebrale Acquisita; Conseguenze; Ictus; Trauma Cranico; Revisione

Autore responsabile per la corrispondenza: Aurora Leontina Lombardi, Dipartimento di Psicologia, Università degli Studi della Campania "Luigi Vanvitelli", Caserta, Italia, e-mail: auroraleontina.lombardi@studenti.unicampania.it

Abstract

Acquired brain injuries (ABIs), including traumatic brain injuries (TBIs) and strokes, frequently result in a wide range of emotional sequelae that profoundly impact patients' quality of life and social functioning. This narrative review examines the emotional consequences that often follow ABIs, including anxiety, depression, apathy, anhedonia, disinhibition, aggression, and deficits in theory of mind. Depression is mainly characterized by pervasive negative subjective feelings and emotional suffering, frequently accompanied by reduced motivation and a tendency to withdraw from daily activities. In contrast, anxiety involves heightened sensitivity to internal and external stimuli, leading to a persistent sense of tension and physiological arousal, which can also affect motivation. While apathy and anhedonia may share some overlap with motivational changes, they are distinct constructs: apathy is defined by emotional indifference and affective detachment, while anhedonia specifically refers to a diminished ability to experience pleasure or interest in previously enjoyable activities. Emotional dysregulation, such as increased irritability, aggression, and impulsive behavior, contributes to interpersonal conflicts and social isolation. Disinhibition can manifest as inappropriate social behaviors, further complicating social reintegration. Additionally, deficits in theory of mind, or the ability to understand others' emotions and perspectives, impair social interactions and relationships. This narrative review aims to highlight the pervasive and multifaceted nature of emotional consequences following ABI. These emotional disturbances are often persistent and challenging, underlying the need for increased awareness and tailored rehabilitation programs, to enhance patients' quality of life.

Keywords

Acquired Brain Injury; Consequences; Stroke; Traumatic Brain Injury; Review

Introduction

Acquired Brain Injury (ABI) refers to any form of brain damage that occurs after birth, excluding injuries related to congenital conditions, degenerative diseases, or birth-related trauma. ABI is categorized in two main types: Traumatic Brain Injury (TBI), caused by external physical forces, and Non-Traumatic Brain Injury (Non-TBI), resulting from internal processes such as strokes, tumors, infections, or anoxia. ABI disrupts neuronal structure and function, leading to impairments that vary in severity and duration, ranging from temporary deficits to lifelong disabilities (Goldman et al., 2022).

ABI is currently one of the leading causes of death and disability among young adults and is considered one of the most prevalent neurological disorders worldwide. Epidemiological data estimate a global incidence of 1 per 500 people, with the highest rates occurring in children under 4, adults under 30, and individuals over 65. A recent study of the Italian population from 2012 to 2021 estimates an ABI rate of 77.3% per 100,000 people. The primary causes include domestic accidents (33.1%) and road accidents (17.7%), predominantly involving males aged 15–24 and the elderly (60+) (Cardile et al., 2024). Individuals affected by ABI often face significant challenges reintegrating into society due to both psychological and physiological consequences, including impairments in motor function, language, cognition, sensory processing, and emotional regulation. Cognitive deficits following ABI often include impairments in arousal, attention, concentration, memory, language, and executive functioning. Memory loss may affect both verbal and nonverbal information, making it challenging to recall conversations, instructions, or visual cues. Executive dysfunction typically involves difficulties in planning, organizing, sequencing, and adapting to new tasks, often accompanied by compromised judgment and impulse control (Rao & Lyketsos, 2000).

Physiologically, ABIs often involve lesions in central pathways that control both simple and complex motor functions. This can result in motor disturbances such as hemiplegia, spasticity, ataxia, and upper motor neuron syndrome, which present with various positive and negative signs affecting both static and dynamic motor skills (Zasler et al., 2012).

Risk factors for ABI are particularly high among males and the elderly, with common mechanisms of injury—including vascular damage, axonal injury, atrophy, and neuronal circuit disruption—shared by both TBI and Non-TBI. Clinical outcomes are influenced by factors such as age, genetics, socioeconomic status, and the specific nature of the injury, which also increases a risk of developing neurodegenerative diseases later in life (Cardile et al., 2024; Goldman et al., 2022). A higher cognitive reserve (CR)—described as the brain's capacity to sustain cognitive performance despite age-related changes or brain injury (Stern et al., 2023)—can mitigate cognitive decline, as it develops through engagement in cognitively stimulating and challenging life experiences (Oliva et al., 2024). Research also indicates that cognitive reserve serves as a protective factor across a variety of neurological conditions, including stroke and traumatic brain injury. These studies underscore the beneficial role of cognitive reserve in improving outcomes across a range of neurodegenerative and injury-related pathologies (Basagni et al., 2023).

Emotional disturbances following ABI are both common and complex, with individuals frequently experiencing disorders such as depression, anxiety, apathy, aggression, and emotional lability. It is essential to distinguish between depression and apathy, as these conditions may initially appear similar and can sometimes be mistakenly classified as the same disorder. The primary point of distinction relies on anhedonia—the inability to experience pleasure—which is a key characteristic of depression but is not necessarily associated with apathy. Apathy, by contrast, is linked to motivational deficits rather than mood disruptions, making it qualitatively different from depression, which typically involves feelings of sadness or hopelessness (Green et al., 2021; Bivona et al., 2019).

Emotional changes following ABI can significantly impact quality of life and present major obstacles to rehabilitation and social reintegration. Although research has historically focused more on cognitive and motor impairments, emotional sequelae have received increasing attention in recent years, due to their critical role in patient outcomes and long-term recovery. Clinicians and healthcare professionals should implement a combination of structured assessments alongside observational methods to effectively identify and monitor emotional and behavioral alterations following ABI. Tools such as the Neuropsychiatric Inventory (NPI, Cummings et al., 1994) and the Personality Assessment Inventory (PAI, Morey et al., 2008) are validated instruments that capture key neurobehavioral and emotional symptoms, including apathy, depression, and aggression. By regularly assessing these symptoms, clinicians can monitor symptom progression over time, allowing for more precise and timely adjustments to treatment plans.

This narrative review aims to summarize the most common emotional sequelae of ABIs, including anxiety, depression, apathy, anhedonia, aggression, theory of mind deficits, and disinhibition.

Anxiety following ABI

Anxiety is a common and debilitating psychological condition that can arise following an ABI. This disorder can significantly affect patients' quality of life, interfere with the rehabilitation process, and further complicate clinical management. However, its prevalence and manifestations can vary significantly due to differences in research methods, diagnostic criteria, and the specific characteristics of the studied samples.

Anxiety is a common and debilitating consequence of TBI and cerebral stroke. Recognizing the associated risk factors and implementing tailored interventions are critical for enhancing patient recovery. Effective management of anxiety following TBI, or stroke requires a comprehensive and individualized approach, where targeted therapies are crucial for improving quality of life and promoting recovery.

Post-TBI Anxiety

TBI is associated with a range of long-term psychological complications, including anxiety disorders. The prevalence of anxiety disorders among TBI patients is notably high. A recent meta-analysis reported that 17.45% of TBI patients exhibit anxiety symptoms, with a relative risk (RR) of 1.9 compared to individuals without TBI (Dehbozorgi et al., 2024). Another meta-analysis indicated that 11% of TBI patients receive a diagnosis of Generalized Anxiety Disorder (GAD), while 37% report clinically significant levels of anxiety (Osborn et al., 2016). These findings underline the significant and persistent prevalence of anxiety in post-TBI patients, with a peak in symptoms occurring between 2 and 5 years after the injury.

The onset of anxiety after TBI is influenced by several factors. A previous history of anxiety disorders significantly increases the risk of developing anxiety post-TBI, with symptoms tending to persist and negatively impact quality of life (Lamontagne et al., 2022). A more specific study examined the pre-existing psychological variables of TBI patients as factors associated with the development of social anxiety. Among the various investigated variables (locus of control, self-esteem, self-efficacy, etc.), the feeling of stigma was identified as an independent predictor of social anxiety (Curvis et al., 2016). According to Mata-Bermudez et al. (2024), alterations in dopamine levels, its receptors, and the DAT transporter, along with changes in the neuronal circuits of brain areas involved in emotional processing, such as the prefrontal cortex, basal ganglia, and amygdala, are observed after a TBI. These alterations may result from both primary brain damage and the neuroinflammatory and metabolic processes that follow the trauma, suggesting that post-TBI depression and anxiety are multifactorial phenomena linked to a combination of dopaminergic dysfunctions and other neurobiological mechanisms.

One of the main challenges in managing post-TBI anxiety is the lack of consistency in the diagnostic methods and criteria across the literature, which complicates the accurate identification and management of these disorders. For example, Osborn et al. (2016) report significant variability in the prevalence of GAD among patients with TBI, with estimates ranging from 11% to 28%. The choice of diagnostic criteria (DSM-IV, DSM-III-R, ICD-10) substantially impacts these prevalence rates, with the ICD-10 yielding a notably lower prevalence (2%) compared to other

systems. Additionally, the use of various diagnostic interviews (e.g., SCID-I, MINI, SADS-L) has further contributed to the variability in findings, with prevalence rates ranging from 2% to 28%.

Post-Stroke Anxiety

Among the psychological complications associated with stroke, anxiety is one of the most common conditions that has a substantial impact on both quality of life and recovery. Studies have shown that approximately 25% of stroke survivors experience acute anxiety, and 20% continue to exhibit anxiety symptoms 3-6 months after the event (Barker-Collo, 2007).

Recent studies have identified factors that influence the onset of post-stroke anxiety disorders, such as age, marital status, and education level. Younger patients and those with higher social deprivation scores tend to report higher levels of anxiety compared to older patients (Broomfield et al., 2015). Pre-existing psychological characteristics, such as neuroticism and pre-existing anxiety, are also significant risk factors for developing post-stroke anxiety and depression (Kootker et al., 2016). However, current studies are often limited by small sample sizes and a lack of homogeneity in diagnostic criteria.

An emerging aspect of post-stroke anxiety research concerns the role of sleep duration. A recent study found that a sleep duration of less than 6 hours per night significantly increases the risk of developing anxiety compared to a sleep duration of more than 7 hours. However, no significant differences were found between genders (Liu et al., 2021). These findings suggest the need for further exploration of how sleep quality and quantity may affect the psychological well-being of post-stroke patients.

Post-stroke anxiety can manifest in various forms, including generalized anxiety, panic attacks, and specific phobias. These psychological issues are a complex aspect of post-stroke recovery, with the primary challenge being the variability in assessment tools. The main tools used in the literature, such as the Beck Anxiety Inventory (BAI) (Beck et al., 1988; Barker-Collo, 2007), the State-Trait Anxiety Inventory (STAI) (Spielberger, 1989), and the Hospital Anxiety and Depression Scale (HADS) (Zigmond & Snaith, 1983; Curvis et al., 2016), assess different dimensions of anxiety but often fail to capture the full spectrum of psychological issues faced by stroke patients. Other tools, such as the Social Phobia Inventory (SPIN) (Connor et al., 2020), focus on specific phobias like social anxiety. However, in these cases as well, the patient's response may be influenced by their neurological condition, complicating the diagnosis. These disorders, in addition to being a response to the trauma experienced, can complicate the recovery process by interfering with rehabilitation and increasing the risk of further complications (Chun et al., 2018). Despite substantial rehabilitative interventions aimed at physical recovery, patients' perceived quality of life does not always improve, indicating that psychological support efforts may be insufficient (Ahlsio et al., 1984).

Treatment options for post-stroke anxiety remain limited and often inadequate to meet patients' specific needs. Some therapies, such as CBT, have shown promising results, but their efficacy must be confirmed by larger and better-designed studies. An investigation by Petrova et al. (2012) examined 198 patients with a first cerebral stroke, finding that 71.2% had affective disorders, including generalized anxiety and phobias. This study highlighted the importance of considering

post-stroke anxiety disorders as relatively independent conditions, with specific risk factors and pathogenesis.

Depression in ABI

Depression is a mood disorder characterized by a persistent sense of sadness and diminished interest in daily activities. The DSM-5 (APA, 2013) classifies depressive disorders into several types: Disruptive Mood Dysregulation Disorder, Major Depressive Disorder, Persistent Depressive Disorder (dysthymia), Premenstrual Dysphoric Disorder, and Depressive Disorder due to Another Medical Condition. All depressive disorders share common symptoms, including feelings of sadness, emptiness, or irritability, along with physical and cognitive changes that significantly affect the individual's ability to function (APA, 2013).

Functional neuroimaging studies have shown that depression is associated with reduced basal metabolism in neocortical structures, particularly the prefrontal and frontal lobes (Sacher et al., 2012), alongside an increase in basal metabolism within the limbic system, including regions like the dorsal raphe nuclei, cingulate cortex, and left prefrontal cortex (Mayberg et al., 1999). Additionally, other studies have identified structural neuroanatomical changes correlated with major depression, such as enlarged lateral ventricles and reduced volumes in areas such as the basal ganglia, thalamus, hippocampus, and frontal lobe (Kempton et al., 2011).

Although depressive disorder diagnoses are established based on specific criteria in the DSM, additional tools are available for identifying and assessing these conditions. Depressive symptoms are usually assessed using tools such as the Beck Depression Inventory-II (BDI-II) and the Montgomery-Åsberg Depression Rating Scale (MADRS). These instruments evaluate key dimensions of depression, with the BDI-II offering a self-reported perspective and the MADRS providing clinician-administered insights. While both tools are widely recognized, their differing emphases—such as the BDI-II's inclusion of somatic and cognitive aspects versus the MADRS's focus on core affective symptoms—may influence reported prevalence rates and severity levels. This highlights the role of measurement choice in shaping study outcomes, particularly in heterogeneous conditions like ABI. (Hudgens et al., 2021; Montgomery et al., 1979).

There is a fair, albeit limited, literature that has explored the relationship between ABI and depression. This is largely due to the fact that early research in this area predominantly focused on the cognitive sequelae of ABI. However, from the 1970s, there has been a shift toward exploring the potential—and unfortunately common—psychopathological consequences of ABI. Among these, depression (diagnosed via the Structured Clinical Interview and the DSM-IV criteria) is one of the most prevalent conditions linked to TBI (Howlett et al., 2022). In a recent article, Iverson and Gardner highlight the well-established link between chronic traumatic encephalopathy (CTE) and various psychiatric disorders, particularly depression. These conditions are often diagnosed, like many affective disorders, using the World Mental Health Survey Initiative Version of the World Health Organization Composite International Diagnostic Interview (WMH-CIDI). This structured diagnostic tool produces diagnoses based on both the International Classification of Diseases, 10th Revision (ICD-10), and the DSM-IV. Their analysis also refers to a series of autopsy studies conducted on populations such as military veterans, professional athletes (including

American football players and boxers). These studies identified widespread histological alterations in the central nervous system, which strongly correlated with the psychiatric symptoms exhibited by these individuals during their lives (Iverson & Gardner, 2021; Omalu et al., 2011; Baugh et al., 2012). Based on these findings, Montenigro and colleagues proposed a classification system for CTE, categorizing it into four types: behavioral/mood variant, cognitive variant, mixed variant, and dementia variant (Montenigro et al., 2014).

Another study by Bryant and colleagues (2010), on the other hand, further corroborates this association, by examining the prevalence of psychiatric disorders among patients admitted to the emergency room following head trauma. Using the Mini International Neuropsychiatric Interview, version 5.5, a brief structured diagnostic interview based on DSM-IV and ICD-10 criteria, their findings indicated that depression was the most common new-onset disorder, occurring in 16.3% of cases. The authors hypothesized that this incidence may be attributed to direct damage to the frontal lobes and subcortical pathways resulting from the injury.

Nevertheless, the study with the largest sample size was conducted by Choi et al., who collected data from the National Health Insurance Service-National Health Information Database (NHIS-NHID) in South Korea. Their results reported an incidence rate of 34.6% of depression among the North Korean population who have experienced head trauma (Choi et al., 2022). They also noted a higher incidence of depression in male patients, along with an apparent correlation between psychiatric symptomatology and the severity of head trauma.

A recent study conducted by Frank et al. (2022) examined the occurrence of depression following stroke, reporting a prevalence rate ranging from 20% to 60%. This finding establishes depression as the most common psychiatric comorbidity in this population. In this context, Feng and colleagues investigated the potential pathogenesis of post-stroke depression, identifying alterations in the frontal regions of the brain and the basal ganglia. However, they emphasized that this topic requires further investigation, as other etiopathological hypotheses suggest a complex interplay involving hyperactivation of the hypothalamic-pituitary axis, neuroinflammation, and immune system dysregulation (Feng et al., 2014).

Apathy following ABI

Apathy is a reduction in goal-directed behaviors and may result from reduced emotional reactivity, interference with motor planning, and failure to self-initiate actions (Levy et al., 2006).

Neuroimaging studies have found associations between apathy and structural and functional alterations in the frontal lobe, particularly in the anterior cingulate gyrus and also in subregions of the basal ganglia (Paul et al., 2005; Roth et al., 2004).

One of the most used tools for assessing apathy in ABIs is the Apathy Evaluation Scale (AES), which measures severity across self-reported (AES-S), informant-reported (AES-I), and clinician-administered formats (AES-C). This tool underscores the importance of considering how measurement tools affect the interpretation of emotional and motivational deficits in ABI population (Marin et al., 1991). The Apathy Scale (AS), a shortened version of the AES scale, is also widely used (Starkstein et al., 1992).

Apathy can also be measured using the Dimensional Apathy Scale (DAS), a multidimensional tool. The DAS was designed to specifically assess three neurobehavioral subtypes of apathy through theoretical analysis and targeted item selection, making it the only scale that comprehensively measures these domains. It detects apathetic deficits related to planning, attention, or organization (Executive subscale), emotion integration (Emotional subscale), and self-generation of behavior or cognitive processes (Initiation subscale) (Radakovic et al., 2016).

Apathy following stroke

The pioneering empirical research on apathy in stroke patients has provided compelling evidence of its existence, showing that it may be more prevalent than previously believed. In this study of Starkstein et al. (1993) apathy was measured using the AS, which was administered by the examiner. They found that approximately 23% of a consecutive series of stroke survivors exhibited signs of apathy. Among those identified as apathetic, nearly half also displayed symptoms of depression. Additionally, this study highlighted that patients with apathy were significantly more likely to experience greater cognitive impairments and deficits in activities of daily living compared to those without apathy. In a study by Brodaty et al. (2005), apathy (measured with the AES) was diagnosed in 26.7% of stroke patients, in contrast to 5.4% in the control group. Subsequently, it has been estimated that the prevalence of apathy (assessed with the AS, using a cut-off score of 12) following a stroke, ranges from 20% to 25%, often correlating with cognitive decline and a chronic trajectory marked by progressive functional deterioration (Jorge et al., 2010). A recent study suggested that apathy after stroke is a common neuropsychiatric consequence, affecting approximately one in three patients. Furthermore, it has been estimated that the prevalence of pure apathy, without concurrent depression, is twice as high as that of pure depression, without accompanying apathy (Caeiro et al., 2013).

Apathy following TBI

In a study conducted by Kant et al. (1998) it was revealed that a significant number of patients with TBI exhibit apathy (assessed with the AES). Furthermore, the authors assert that apathy can occur independently of depression, suggesting that its underlying pathophysiological mechanisms may differ from those associated with depression.

Similarly, a study by Ciurli et al. (2011) confirmed that apathy (measured with the NPI), is the most prevalent neuropsychiatric symptom among subjects with TBI, identified in 42% of cases. Within the sample of participants with severe TBI, apathy was found to be the most common neuropsychiatric disorder. Consequently, it can be stated that apathy is a prevalent behavioral complication in TBI patients, affecting at least half of individuals at some stage during the post-TBI recovery period. While apathy in TBI is frequently associated with both depression and cognitive deficits, it can also manifest as an independent condition in numerous cases (Starkstein et al., 2014).

Apathy following ABI

In patients with subcortical or hypoxic brain damage, as well as those with unilateral right hemisphere damage, apathy, assessed with the AES, has been found to be most severe as a result of ABI. The strong association between apathy and the location of the lesion suggests that the latter arises directly from the lesion and not from secondary psychological or socio-environmental problems. This finding points to an underlying dysfunction in the brain's motivational systems, supporting the classification of apathy in patients with acquired brain damage as a disorder of organic origin (Andersson et al., 1999).

This finding points to an underlying dysfunction in the brain's motivational systems, supporting the classification of apathy in patients with ABIs as a disorder of organic origin (Andersson et al., 1999).

Anhedonia and ABI

Anhedonia is defined as the inability to experience pleasure in activities that are usually considered enjoyable, such as physical exercise, hobbies, sexual activities, and social interactions, especially when compared to similar experiences that were previously perceived as pleasurable (Calabrò et al., 2012). According to the DSM-5, anhedonia manifests as a "markedly diminished interest or pleasure in all, or almost all, activities most of the day, nearly every day" (APA, 2013). On the other hand, the International Classification of Diseases (ICD-10) by the World Health Organization does not use the term "anhedonia" but refers to "loss of interest and pleasure" (World Health Organization, 2004).

In clinical practice, the term "anhedonia" encompasses a complex and multifaceted dysfunction related to the reward system, which becomes more evident compared to healthy subjects. However, a study by Ho and Sommers (2013) found that about 45% of studies measuring anhedonia do not provide a clear definition of the concept. The ability to experience pleasure is not limited to desire, reinforcement, and subjective pleasure but also includes cognitive aspects such as the ability to anticipate and predict rewards, the memory associated with these experiences, the association between value and relative cost, the effort required to obtain rewards, the integration of this information, the resulting decision-making process, and the self-motivation to engage in goal-directed behaviors (Der-Avakian & Markou, 2012). Consequently, reward processing is strongly influenced by executive functions (Berkman et al., 2011; Berkman et al., 2014).

Some authors have proposed a distinction between consummatory and anticipatory anhedonia to better describe the lack of a specific aspect of the pleasure process. Consummatory pleasure refers to the enjoyment felt during the execution of a pleasurable activity, whereas anticipatory pleasure relates to the expectation of pleasure deriving from future experiences. These concepts can explain the different behavioral, cognitive, and emotional experiences related to anhedonia. Other authors emphasize the need to distinguish between consummatory and motivational anhedonia, focusing on decision-making processes (Treadway & Zald, 2011).

From a neurobiological perspective, anticipatory pleasure is strongly, though not exclusively, correlated with the mesolimbic dopamine pathway, while the serotonergic and opioid systems

appear to be more centrally involved in consummatory pleasure (Berridge & Robinson, 1998; Wise, 2008). Additionally, anticipatory pleasure is linked to motivational processes that promote behaviors aimed at achieving desired rewards (Schultz, 2002; Carver, 2001; Gard et al., 2007).

Anhedonia is linked to reduced reactivity and connectivity in neural networks involved in the perception and evaluation of rewarding or pleasurable stimuli, including the orbitofrontal cortex and ventral striatum/nucleus accumbens. It also involves disruptions in areas responsible for cost-benefit analysis and decision-making, such as the anterior cingulate cortex, ventromedial prefrontal cortex, and dorsolateral prefrontal cortex. Additionally, anhedonia affects reward processing and memory integration in the mesolimbic pathway (including ventral striatum/nucleus accumbens, ventral tegmental area/substantia nigra, and hippocampal connections), as well as salience attribution within the amygdala, and related limbic and paralimbic structures (Preda, 2014).

As the conceptualization of anhedonia has expanded, assessment tools have been developed to measure dimensions, such as anticipatory and consummatory pleasure, motivation, and effort across different reward domains. For instance, the Temporal Experience of Pleasure Scale (TEPS) (Gard et al., 2006) distinguishes between anticipatory and consummatory pleasure, while the Motivation and Pleasure Scale - Self-Report (MAP-SR), focuses on aspects of anticipation, motivation, effort and pleasure, mainly in the social domain (Llerena et al., 2013). Tools like the Specific Loss of Interest and Pleasure Scale (SLIPS) (Winer et al., 2014) and the Dimensional Anhedonia Rating Scale (DARS) (Rizvi et al., 2015) assess broader aspects of interest, motivation, effort, and pleasure. Similarly, the Positive Valence Systems Scale (PVSS) evaluates reward-related aspects specified in the positive valence systems (PVS) framework within the Research Domain Criteria (RDoC) (Khazanov et al., 2020). While these tools provide valuable insights into different facets of anhedonia, their varied emphasis on specific domains (e.g., physical versus social rewards) may influence how anhedonia is characterized and its prevalence. This underscores the importance of aligning tool selection with study objectives to ensure accurate interpretation of results.

Given the complexity of anhedonia, it is considered a symptom of various neuropsychiatric conditions, including depression (Pelizza & Ferrari, 2009), bipolar disorder (Tso et al., 2014), and ABIs of varying severity (Rao et al., 2007).

Anhedonia is a transdiagnostic symptom, in that it extends beyond major depression to social anxiety disorder and generalized anxiety disorder (Kashdan et al., 2011), as well as schizophrenia (Watson and Naragon-Gainey 2010) and substance abuse (Thomsen et al., 2015).

Anhedonia, assessed via low positive emotionality, has been shown to prospectively predict the development of both depression and anxiety, even when controlling for baseline symptoms (Kendall et al., 2015; Khazanov & Ruscio, 2016). Once disorders emerge, anhedonia serves as a strong predictor of a more unfavorable long-term course of major depression (Morris et al. 2009). It also forecasts poor psychosocial functioning following improvements in depressive mood (Vinckier et al., 2017) and increases the likelihood of depression recurrence (Wichers et al., 2010). Furthermore, anhedonia is a significant predictor of suicidal ideation and behavior (Winer et al., 2014; Ducasse et al., 2018), even after accounting for other cognitive and emotional symptoms of depression (Fawcett et al., 1990; Ballard et al., 2017) as well as other risk factors such as history of suicide attempts, childhood trauma, marital status, gender, and age (Ducasse et al., 2020).

Anhedonia and Stroke

Motor, cognitive, and language deficits are common post-stroke consequences that limit the ability to perform daily activities, reducing participation in family, social, work, and community life (Hartman-Maeir et al., 2007; Tang et al., 2020). The reduction in functional autonomy in stroke survivors negatively affects their emotional well-being and quality of life, contributing to the onset of mood disorders (Patel et al., 2002; Marcheschi et al., 2018; Diamond et al., 2023). Specifically, it is estimated that between 25% and 79% of stroke patients suffer from post-stroke depression (PSD), a condition linked to the physical and psychological adversities they face (Whyte & Mulsant, 2002). Previous studies have shown that PSD has a negative impact on motor and cognitive recovery, contributing to a 15% increase in overall disability (Paolucci et al., 2019). This detrimental effect has been associated with deficits in the regulation of motivational behaviors (Gainotti et al., 2001).

Although many studies have focused on the negative impact of apathy on post-stroke rehabilitation (Marin, 1991; Sibon et al., 2012), motivational mechanisms and goal-directed behaviors can also be impaired by anhedonia, a condition still understudied in the stroke population. In this context, the disruption of dopaminergic networks following a stroke could trigger mood disorders and cognitive deficits, including memory dysfunction, contributing to the development of anhedonia (Piamarta et al., 2004; Terroni et al., 2015). Considering the crucial role of anhedonia in the motivational aspects of post-stroke recovery, a recent study by Ashaie et al. (2023) explored the association between three dimensions of PSD somatic symptoms, negative affect, and anhedonia. The results showed that anhedonia predicted an increase in somatic symptoms during the first year after discharge from rehabilitation, indicating its impact on physical distress rather than the direct effects of the stroke. Consequently, the authors emphasized the importance of examining specific dimensions of PSD, such as anhedonia, to better understand its development and etiology, allowing for targeted treatments and improved therapeutic outcomes (Ashaie et al., 2023).

A recent cohort study conducted by Segura et al. (2024) revealed a significantly higher prevalence of anhedonia among stroke patients (18.5-19.7%) compared to healthy controls (4.3%). Particularly, the study found that the levels of anhedonia among the patients were consistent across various factors, including stroke etiology, time since the event, affected hemisphere, lesion location, and motor and cognitive function levels. Additionally, anhedonia was associated with lack of motivation and increased negative emotional states, such as fatigue and anger, which adversely affected emotional well-being and quality of life, potentially leading to decreased participation in rehabilitation programs. Other studies have emphasized the role of anhedonia as a predictor of somatic symptoms of depression, such as fatigue, within the first year following rehabilitation discharge (Ashaie et al., 2023). It may also contribute to a self-perpetuating cycle of behavioral deterioration (Mallorqui et al., 2022). Furthermore, the literature indicates that older stroke patients face a heightened risk of experiencing long-term anhedonia. This increased vulnerability is attributed not only to a greater likelihood of social isolation (Yeh & Lo, 2004) but also to the neurophysiological damage resulting from brain injury.

Anhedonia and TBI

From a neuropathological perspective, TBI is characterized by changes in various cortical areas, subcortical structures, and the white matter tracts that connect them. Recent evidence suggests that diffuse neuronal damage and cell loss can progress over weeks or months after the initial insult in selectively vulnerable regions of the prefrontal cortex, hippocampus, thalamus, striatum, amygdala, and forebrain nuclei. The resulting functional changes in neuronal circuits may constitute the neurological substrate of the cognitive and behavioral deficits frequently observed following TBI (Bigler et al., 2002; McIntosh et al., 1998).

Psychopathological disorders, particularly mood disorders, occur in the context of profound changes in cognitive and emotional processing following TBI. There is extensive literature regarding the cognitive changes observed after TBI. Depression occurs in approximately 25% of TBI patients (Fedoroff et al., 1992; Jorge et al., 1993). Feelings of loss, demoralization, and discouragement observed shortly after the injury are often followed by symptoms of persistent dysphoria. Fatigue, irritability, suicidal thoughts, anhedonia, lack of interest, and insomnia are observed in a considerable number of patients 6-24 months or even longer after TBI (Hinkeldy et al., 1990; Van Zomeran et al., 1985). In TBI patients, anhedonia reduces quality of life and may hinder rehabilitation (Jorge et al., 2005).

In a retrospective study (Lewis et al., 2015), data from 192 TBI participants from Phase III of the Vietnam Head Injury Study, a long-term follow-up study on veterans primarily with focal penetrating TBI (Raymont et al., 2011), were analyzed. Participants underwent neurological and psychiatric examinations, neuropsychological testing, and brain CT scans. The study aimed to examine the expression of anhedonic symptoms related to localized TBI. The primary finding was that anhedonic symptoms were more pronounced in individuals with damage to the right ventrolateral prefrontal region compared to those with injuries in other regions.

Disinhibition after ABI

The term “disinhibition syndrome” describes a wide range of behaviors that are inappropriate to the context and reflect the inability to control and repress some reactions. Different symptoms have as their common mechanism the lack of inhibition, such as impulsivity, inappropriate behaviors, aggressivity, and what Eslinger and Damasio (1985) called “acquired sociopathy”, which is the loss of sensitivity, ability to respect social rules, and the account of consequences of aggressive actions on other people. The definition of disinhibition provided by Arciniegas and Wortzel (2014) is “socially and contextually inappropriate nonaggressive verbal, physical, and sexual acts that reflect a lessening or loss of inhibition and/or inability to appreciate social or cultural behaviors norms”. Patients with disinhibition may have a normal level of cognitive intelligence and compromised emotional intelligence (Bar-On et al. 2003). According to the hypothesis of “somatic markers” (Damasio, 1994) a poor detection of somatic signals is the core feature that leads to disadvantageous choices for the people and for the way they relate to others. Blair and Cipollotti (2000) suggested an important mechanism for social cognition and control of aggressivity: the Social Response Learning. A lesion of the orbito-frontal cortex damages this system and can cause disinhibition and acquired sociopathy. This finding has been confirmed by

recent studies using modern technologies like MRI. For example, Knutson et al. (2015) conducted a whole-brain analysis, and identified several brain regions involved in behavioral disinhibition, including the right orbitofrontal regions, bilateral insula, right temporal lobe, left frontal, precentral and postcentral regions, and bilateral gyrus rectus.

Further evidence comes from He et al. (2020), who analyzed structural MRI data from 361 healthy adults as part of the Human Connectome Project (HCP). They used the Connectome - based predictive model (CPM) to identify brain areas that are associated with inhibitory control. Their findings revealed a “high inhibitory control network” involving nodes in the prefrontal, orbitofrontal cortex, and inferior frontal gyrus, as well as a “low inhibitory control network”, with nodes distributed across parietal, occipital, and limbic areas.

In a more recent study, Jenkins et al. (2024), found that disinhibition correlates with reduced global efficiency of Cognitive Control Network, and its connection with the Salience Network. Increasingly, studies of neural connectivity in disinhibition and other neuropsychiatric disorders are finding that these conditions are not the result of dysfunction in a single brain area. Instead, they may be better explained by widespread disconnection across distant brain regions.

Disinhibition in stroke.

Multiple studies indicate that between 5% and 76% of stroke patients show disinhibition at various time points ranging from 4 days to 4 years post-stroke (van Almenkerk et al., 2012; Angelelli et al., 2004; Aybek et al. 2005; Ghika-Schmid et al., 1999; Greenop et al. 2009; Mok et al., 2010; Buijck et al., 2012; Wong et al. 2014). However, these data are unreliable due to the frequent underdiagnosis and undertreatment of disinhibition. This presents a significant issue, as untreated disinhibition can result in prolonged stays in rehabilitation facilities and a slower recovery (Buijck et al., 2012). Ito et al. (2023) showed that patients with a MMSE score lower than 18 had higher Neuropsychiatric Inventory-Questionnaire Severity scores in disinhibition and Neuropsychiatric Inventory-Questionnaire Distress scores. This result shows that the severity and frequency of disinhibition is significantly higher in Subacute Stroke patients with severe cognitive impairment than in patients with mild cognitive impairment, while depression and apathy occur regardless of the severity of cognitive impairment. This study suggested that cognitive, behavioral and psychological symptoms should be early managed in stroke patients with cognitive impairment.

Disinhibition in TBI

There is considerable variability in studies regarding the prevalence of disinhibition among patients with TBI. Warriner et al. (2003), using the Minnesota Multiphasic Personality Inventory, reported that only 13% of TBI patients showed disinhibition, while Kelly et al. (2008) assessed the disinhibition through the Overt Behavior Scale (OBS, Kelly et al., 2006), and found a prevalence of 85.8%. This lack of agreement may be due to inconsistent definitions of the construct, the different assessment tools used, and variations in the severity of injuries within the samples (Osborne-Crowley and McDonald, 2018). Nonetheless, many studies have showed that disinhibition in TBI patients can lead to various long-term consequences, including caregiver

burden (Brooks et al., 1986), family dysfunction (Douglas & Spellacy, 1996), reduced quality of life for patients (Koskinen, 1998), challenges in social reintegration (Simpson et al., 1999), and an increased risk of suicidal thoughts (Juengst et al., 2014).

Aggression after ABI

Aggression following ABI is typically characterized by disproportionate or inappropriate behaviors, which may be verbal or physical, and can be directed toward others, objects, or oneself. The manifestation of aggression can vary considerably; it depends on numerous factors, including the location and extent of the injury, as well as the individual's pre-injury personality (Neumann et al., 2015; Rao et al. 2009). Additionally, other co-occurring factors, such as cognitive impairment, emotional dysregulation, and environmental stressors, may exacerbate the aggressive behaviors. However, it remains unclear whether the varying expressions of aggression are to be considered as distinct syndromes or if they could be better understood as part of a continuum of symptoms (Neumann et al. 2015; Rao et al., 2009). Aggression following ABI can adversely impact family dynamics, social relationships, and community reintegration; furthermore, it can exacerbate the burden on families and may lead to the exclusion of the patient from treatment programs, thereby perpetuating a cycle of further isolation and diminishing access to essential social support systems (Hart et al., 2017). The *Overt Aggression Scale – Modified for Neurorehabilitation* (OAS-MNR) (Alderman et al., 1997) was developed as a standardized tool to assess aggression in neurorehabilitation settings using an operant framework. It categorizes aggression into four types—verbal aggression, and aggression toward objects, self, and others—each rated on a severity scale from mild to very severe based on observable behaviors. The OAS-MNR reliably records details of aggression episodes, including context and antecedents, and has proven useful in clinical practice, research, and program evaluation; it is one of the only tools, along with its extended version (Giles & Mohr, 2007), that has been specifically validated for individuals with ABI.

A variant, the *St Andrew's Sexual Behavior Assessment* (SASBA), assesses potentially inappropriate sexual behaviors. Both OAS-MNR and SASBA are considered "focal" measures, providing in-depth insights into specific behaviors. However, in cases where tracking every incident is impractical, broader "global" assessment tools, like the *Overt Behavior Scale* (OBS) and the *St Andrew's–Swansea Neurobehavioral Scale* (SASNOS), are valuable. These tools measure challenging behaviors, including aggression, in community settings for ABI populations. The OBS and SASNOS are easy to administer, providing a profile of strengths and weaknesses to support rehabilitation planning (Alderman et al., 2013).

The choice of tool can influence findings, with focal measures like the OAS-MNR capturing detailed episodic data, while global tools like the OBS and SASNOS provide a comprehensive behavioral overview. Selecting the appropriate tool is thus critical for aligning measurement with study objectives and ensuring accurate characterization of aggression in ABI populations.

Aggression after stroke

During the acute stage, stroke patients may exhibit aggressive behaviors, both physical and verbal, such as kicking, biting, pushing, throwing objects, screaming, cursing, and shouting.

However, also in the subacute stage, patients often become more irritable, impulsive, and hostile, frequently displaying uncontrollable anger towards family members over minor issues. These symptoms are commonly referred to as post-stroke anger proneness (PSAP) (Kim, 2016). Such disruptive behavior during treatment or rehabilitation can cause significant distress to caregivers and result in communication breakdowns between patients and healthcare professionals, thereby hindering and delaying the rehabilitation and recovery process (Choi-Kwon et al., 2022).

The prevalence of aggressive behavior within the first week following a stroke is reported to range from 17% to 23%, increasing up to 32% between 3- and 12-months post-stroke. Aggression in stroke patients has been found to be correlated with factors such as younger age, a history of depression, and speech difficulties, (i.e., dysarthria). Additionally, it is often associated with concurrent depressive symptoms, impaired cognitive function, and motor dysfunction. (Choi-Kwon et al., 2022; Lau et al., 2017).

Numerous studies have attempted to establish a relationship between the locations of stroke lesions and PSAP; however, the findings have been inconsistent (Choi-Kwon et al., 2022). For instance, one study utilizing CT scans suggested that lesions in the left hemisphere and those situated closer to the prefrontal cortex were associated with aggressive behavior; however, they used a small sample size, which limits the reliability of this finding (Paradiso et al., 1996). In contrast, a study, conducted by Kim et al. using MRI, highlighted that PSAP was strongly linked to lesions in the fronto-lenticulo-capsular-pontine base area, while patients with lesions in other brain regions rarely exhibited PSAP (Kim et al., 2002). Conversely, other studies have reported no significant correlation between the location of stroke lesions and the occurrence of PSAP (Santos et al., 2005). Additional studies have indicated that PSAP may be linked to personality changes associated with frontal lobe lesions, which can lead to increased aggression or the amplification of preexisting aggressive tendencies. Furthermore, alongside the relationship between cognitive impairment and PSAP, patients may experience anger as a result of frustration related to impaired communication abilities, inappropriate responses to stimuli (such as pain or noise), or negative attributions regarding other's behaviors (Choi-Kwon et al., 2022; Neumann et al., 2015).

Aggression following TBI

Aggression is one of the most common consequences after TBI, with prevalence estimates ranging from 11% to 34% (Rao et al., 2009). The variation in reported prevalence rates across studies can be attributed to differences in study design, duration, population characteristics, and the criteria used to define and categorize aggression. Moreover, the complexity of post-TBI aggression, which may also be a symptom of delirium, mood disorders, or personality changes secondary to TBI, further complicates the ability to generalize findings (Visscher et al., 2010; Rao et al., 2009). Verbal aggression is typically the most prevalent subtype, in both the acute and the chronic settings (Roy et al., 2017).

Aggression is considered problematic following TBI due to the frequency, intensity, and apparent inability to regulate negative emotions, such as anger, and related behaviors. These emotional deficits are commonly described as being unpredictable, irrational, inappropriate, and disproportionate to the circumstances (Neumann et al., 2015). In fact, aggressive behaviors not

only disrupt rehabilitation efforts but also impose a significant burden on both patients and caregivers. A comprehensive understanding of the correlates and predictors of aggression is crucial for effective prevention and treatment strategies, which could ultimately enhance rehabilitation outcomes during the critical early post-TBI period (Rao et al., 2009; Visscher et al., 2010).

In acute rehabilitation, aggressive behavior has been linked to a range of factors, including impaired language function, symptoms of post-traumatic stress disorder, frontal lobe injuries, a history of premorbid psychosocial issues and substance abuse, delirium, mood disorders (i.e., depression) and psychosis, late-onset partial complex epilepsy, a higher incidence of medical comorbidities, disorientation to place and time, and the use of anticonvulsant medications (Baguley et al., 2006).

The neuroanatomical mechanism underlying post-TBI aggression remains unclear, but it could be the result of an imbalance between inhibitory pathways in the frontal cortex and limbic structures that regulate impulse (Kim, 2002). For instance, in a study conducted by Tateno et al. (2003) frontal lesions were found to be associated with aggression; this would suggest that frontal lobe injury can cause damage to the ascending serotonergic pathways, which can contribute to the pathophysiology of both depression and aggressive behavior (Rao et al., 2009; Tateno et al., 2003).

Theory of mind and acquired brain injury

The term Theory of Mind (ToM) refers to the process of mentalizing, which is fundamental for social cognition and social interactions (Clausi et al., 2019). This mechanism is distinctly different from emotional contagion, an automatic process involving the recognition of others' emotions; ToM involves more complex processes, including the ability to attribute mental states different from one's own to others and to adopt another person's perspective in order to understand and predict their behavior. Conceptually, a distinction exists between "hot" functions, such as processing emotions, and "cold" functions such as understanding another's perspective. Among the hot functions, affective empathy refers to the ability to connect emotionally with the feelings of others while recognizing that these feelings are separate from our own. Emotion perception, another hot function, involves interpreting a person's emotional state based on cues like facial expressions or tone of voice. Cold functions, on the other hand, involve skills such as explaining the behavior of others by understanding their thoughts, appreciating others' points of view, and interpreting pragmatic inferences (McDonald, 2013).

It has been shown that patients with brain injuries may exhibit reduced empathy (Shamay-Tsoory et al., 2004), social isolation (Lezak et al., 2004), difficulties in understanding irony or sarcasm (Martin & McDonald, 2005; Channon et al., 2005), and poor social skills (Spatt et al., 1997). These factors can contribute to challenges in social reintegration.

Patients with brain lesions tend to fail tasks that require reasoning about mental states, making both ToM and mentalizing tasks predictors of deficits in social behavior. Various tasks are used to assess ToM, generally categorized into "first-order" tasks adapted from developmental psychology (e.g. the Sally-Anne test) and "second-order" tasks, with the "Faux Pas" test (Stone et al., 1998) being the most frequently used. This test involves presenting stories in which a character makes a faux pas, such as saying something inappropriate to the listener. To recognize the faux pas, the

subject must understand that the protagonist of the story is unaware of making a social mistake (Clausi et al., 2019). The Sally-Anne test assesses understanding of false beliefs by asking participants to predict where a character will look for an object based on their perspective, rather than reality (Korkiakangas et al., 2016). (Korkiakangas et al., 2016). In the faux pas test, subjects are presented with stories where a character unintentionally says something socially inappropriate, requiring participants to recognize the social mistake and understand the protagonist's lack of awareness (Clausi et al., 2019). Participants answer questions designed to evaluate their ability to detect the faux pas, attribute intentions, and assess others' emotional states (Söderstrand & Almkvist, 2012). These tasks provide valuable insights into ToM impairments, particularly in populations with ABI, by assessing key cognitive and affective dimensions of social cognition. However, differences in task complexity, focus on verbal versus non-verbal cues, and reliance on executive functioning may influence findings, underscoring the importance of tool selection in capturing the full spectrum of ToM abilities.

A review by Abu-Akel & Tsoory (2010) highlighted that both cortical and subcortical regions form distinct, yet interacting, networks that support the cognitive and affective representation of mental states, both in relation to oneself and to others. The cognitive network involves the dorsomedial prefrontal cortex, the dorsal anterior cingulate cortex, and the dorsal striatum. The affective network involves the ventromedial prefrontal and orbitofrontal cortex, the ventral part of the anterior cingulate cortex, the amygdala, and the ventral striatum. Additionally, attention and selection systems, particularly the parietal-temporal junction, play a critical role in facilitating these processes. Neurochemically, the dopaminergic and serotonergic systems seem to play key roles in these functions.

ToM and TBI

The relationship between ToM and TBI has been extensively examined in the literature, consistently revealing deficits in ToM tasks among individuals with TBI. Most studies have used narrative stories (Bibby & McDonald, 2005; Martin & McDonald, 2005) or videos (McDonald & Flanagan, 2004; Turkstra et al., 2004) to assess ToM. However, Henry et al. (2006) adopted a less cognitively demanding test to minimize the influence of cognitive load. Despite the reduction in task complexity, TBI patients still showed considerable impairments, scoring significantly lower than the control group.

These findings align with broader research on the relationship between ToM and executive functioning. For instance, Apperly et al. (2007) demonstrated that both ToM tasks, (e.g. false belief tasks) and non-ToM tasks, such as "false photograph" tasks—where perspective-taking is not required—show deficits in individuals with brain damage. This suggests that, despite arguments for the specificity of ToM as a cognitive process (Frith & Frith, 2003; Saxe et al., 2004), efficient executive functioning is essential for performing mentalizing tasks. In the study by Apperly et al. (2007), impaired performance on mentalizing tasks correlated with deficits in a phonemic fluency task, a measure of executive functioning. This correlation was not found with a purely emotional task that required subjects to identify the emotion expressed on a face, although performance on this task was also impaired compared to controls. Overall, in addition to social cognition tasks,

individuals with TBI show below-average performance in tasks assessing both emotional and cognitive empathy (de Sousa et al., 2010).

All these difficulties—in experiencing empathy, understanding others' mental states, and pure executive functioning—constitute significant barriers to full recovery after a brain injury and hinder the return to premorbid social life.

ToM and stroke.

Stroke has also been widely investigated in relation to ToM. Research in this area has predominantly examined the differences between lesions in the two hemispheres and their impact on both the cognitive (beliefs about beliefs) and emotional (beliefs about feelings) components of ToM. A study conducted by Yeh and Tsai (2014) compared the performance of subjects with right hemisphere lesions to those with left hemisphere lesions and compared these results with a control group of healthy individuals. The study involved tasks that assessed both cognitive and affective ToM in verbal (*faux pas*) and non-verbal modalities. The underlying hypothesis of this study primarily concerned empathy, suggesting that deficits in affective tasks are rooted in reduced empathic capacity caused by the lesion; therefore, tasks assessing empathy were also employed. The results showed that subjects with right hemisphere lesions exhibited poorer performance in both non-verbal cognitive ToM tasks and cognitive empathy tests compared to subjects with left hemisphere strokes. However, both groups were impaired compared to controls in both affective and cognitive ToM tasks.

Similar to what has been discussed for individuals with TBI, stroke patients also show poor performance on ToM tasks that correlate with tests assessing executive functioning, such as the Brixton Test (Hamilton et al., 2017). Therefore, it is crucial to consider the decline in cognitive flexibility, which may affect mentalizing tasks and, more broadly, the overall psychosocial functioning of individuals with brain lesions.

All these findings taken together, including those related to TBI and stroke, highlight the importance of administering tasks that assess ToM in cases of ABI. This approach is vital for developing rehabilitation programs that support social reintegration and promote the individual's complete recovery.

Therapeutic approaches and future directions

Improving the quality of life for patients with emotional sequelae following ABI requires a wide range of therapeutic approaches, yet achieving a clear consensus on optimal treatment strategies remains challenging. There is growing emphasis on multidisciplinary care, with an increase in advocating for a team-based approach that includes traditional healthcare providers, such as psychiatrists and neurologists, as well as specialists like occupational therapists, neuropsychologists, and psychotherapists. Such frameworks extend beyond hospital settings, actively engaging with community resources to provide comprehensive care.

Treatment options for post-stroke anxiety remain limited and often inadequate to meet patients' specific needs. While CBT has shown promise in reducing anxiety symptoms, further research is needed to confirm its efficacy in this population. Petrova et al. (2012) studied 198 patients with a

first cerebral stroke and found that 71.2% experienced affective disorders, including generalized anxiety and phobias. Post-TBI anxiety is typically managed through both pharmacological and psychotherapeutic approaches. Studies have shown CBT to be effective in alleviating post-TBI anxiety and depression, with benefits lasting up to three months (Barua et al., 2024). However, pharmacotherapy remains the most common intervention, especially in cases of post-traumatic stress disorder, though mindfulness has proven helpful in youth with TBI, reducing anxiety and enhancing psychological functioning (Soo et al., 2019).

For patients with depression, cognitive behavioral therapy (CBT) has shown mixed efficacy, while repetitive transcranial magnetic stimulation (rTMS) has produced promising outcomes (Frank et al., 2022). Addressing apathy often includes pharmacological treatments like selective serotonin reuptake inhibitors (SSRIs), though these may only be effective when depressive symptoms coexist, and certain antidepressants may even impair motivation. Rehabilitative techniques, such as neuropsychological counseling, focus on restoring goal-directed behaviors through goal-setting, future planning, and self-monitoring, often complemented by CBT. Dopaminergic pharmacotherapy and rTMS have also shown some potential in treating apathy (Tay, 2021).

Treatment options for anhedonia often focus on reducing negative affect, but current psychotherapies, including CBT and mindfulness-based cognitive therapy, have limited impact on enhancing positive affect (Boumparis et al. 2016). Traditional therapies like cognitive therapy and antidepressants improve negative affect but only moderately increase positive affect levels, which remain below those in the general population. New approaches targeting positive valence systems show promise for effectively addressing anhedonia (Dunn et al., 2020; Sandman, 2022).

For aggression management following ABI, an integrative, multidisciplinary strategy is essential, starting with a thorough assessment. While pharmacological intervention may be required, particularly in acute settings, a more integrative approach should be adopted, incorporating cognitive-behavioral interventions tailored to individual needs. Disinhibition rehabilitation may benefit from behavioral management and self-management strategies, as well as electrical aversion therapy, though CBT alone has shown limited effectiveness in reducing disinhibition symptoms (Verberne et al., 2019).

Rehabilitation for ToM deficits can draw on methods used with children, such as conversation-based training. In this approach, participants discuss stories that highlight social nuances under the guidance of a trainer, who encourages perspective-taking through structured feedback and group discussion, followed by individualized exercises (Lecce et al., 2014; Caputi et al., 2020).

Overall, while traditional pharmacological and behavioral therapies continue to be widely used, the most promising outcomes may arise from a comprehensive, interdisciplinary approach that utilizes community resources and personalized care plans to address the complex needs of these patients.

Conclusions

This narrative review summarized the most common emotional sequelae after ABI such as depression, anxiety, aggression, anhedonia, apathy, disinhibition, and deficit in theory of mind.

These emotional changes can arise from both direct neurological damage and the psychological stress of coping with the aftermath of the injury.

Despite the prevalence of emotional sequelae, it is only in recent times that growing attention has been given to improving early detection and intervention. The integration of neuropsychological assessments and interdisciplinary rehabilitation programs have been found to improve emotional outcomes. Therefore, tailored therapeutic strategies, including psychotherapy, pharmacological interventions, and cognitive rehabilitation, are vital for addressing both the emotional and cognitive consequences of ABI.

In conclusion, emotional changes following ABI can be pervasive and long-lasting, demanding more attention from clinicians and researchers. Future studies should focus on the long-term trajectory of these emotional sequelae and the development of more effective, individualized interventions to enhance recovery and improve patients' quality of life.

References

- Abu-Akel, A., & Shamay-Tsoory, S. (2011). Neuroanatomical and neurochemical bases of theory of mind. *Neuropsychologia*, *49*(11), 2971-2984.
- Ahlsio, B., Britton, M., Murray, V., & Theorell, T. (1984). Disablement and quality of life after stroke. *Stroke*, *15*(6), 886-890.
- Alderman, N., Knight, C., & Brooks, J. (2013). Rehabilitation Approaches to the Management of Aggressive Behaviour Disorders after Acquired Brain Injury. *Brain Impairment*, *14*(1), 5-20.
- Alderman, N., Knight, C., & Morgan, C. (1997). Use of a modified version of the Overt Aggression Scale in the measurement and assessment of aggressive behaviors following brain injury. *Brain Injury*, *11*, 503-523.
- American Psychiatric Association. (2013) Diagnostic and Statistical Manual of Mental Disorders (5th ed.). Arlington: American Psychiatric Organization.
- Andersson, S., Krogstad, J. M., & Finset, A. (1999). Apathy and depressed mood in acquired brain damage: relationship to lesion localization and psychophysiological reactivity. *Psychological Medicine*, *29*(2), 447-456.
- Angelelli, P., Paolucci, S., Bivona, U., Piccardi, L., Ciurli, P., Cantagallo, A., Antonucci, G., Fasotti, L., Di Santantonio, A., Grasso, M. G., & Pizzamiglio, L. (2004). Development of neuropsychiatric symptoms in poststroke patients: a cross-sectional study. *Acta psychiatrica Scandinavica*, *110*(1), 55-63.
- Apperly, I. A., Samson, D., Chiavarino, C., Bickerton, W. L., & Humphreys, G. W. (2007). Testing the domain-specificity of a theory of mind deficit in brain-injured patients: Evidence for consistent performance on non-verbal, "reality-unknown" false belief and false photograph tasks. *Cognition*, *103*(2), 300-321.
- Arciniegas, D. B., & Wortzel, H. S. (2014). Emotional and behavioral dyscontrol after traumatic brain injury. *The Psychiatric clinics of North America*, *37*(1), 31-53.
- Ashaie, S. A., Funkhouser, C. J., Jabbarinejad, R., Cherney, L. R., & Shankman, S. A. (2023). Longitudinal Trajectories of Post-Stroke Depression Symptom Subgroups. *Neurorehabilitation and neural repair*, *37*(1), 46-52.
- Aybek, S., Carota, A., Ghika-Schmid, F., Berney, A., Melle, G. V., Guex, P., & Bogousslavsky, J. (2005). Emotional behavior in acute stroke: the Lausanne emotion in stroke study. *Cognitive and behavioral neurology: official journal of the Society for Behavioral and Cognitive Neurology*, *18*(1), 37-44.
- Baguley, I. J., Cooper, J., & Felmingham, K. (2006). Aggressive behavior following traumatic brain injury: how common is common?. *The Journal of head trauma rehabilitation*, *21*(1), 45-56.
- Barker-Collo, S. L. (2007). Depression and anxiety 3 months post stroke: Prevalence and correlates. *Archives of Clinical Neuropsychology*, *22*(5), 519-531.
- Bar-On, R., Tranel, D., Denburg, N. L., & Bechara, A. (2003). Exploring the neurological substrate of emotional and social intelligence. *Brain: a journal of neurology*, *126*(Pt 8), 1790-1800.
- Barua, U., Ahrens, J., Shao, R., MacKenzie, H., Wolfe, D., Sequeira, K., ... Mehta, S. (2024). Cognitive behavioral therapy for managing depressive and anxiety symptoms after brain injury: A meta-analysis. *Brain Injury*, *38*(3), 227-240.
- Basagni, B., Di Rosa, E., Bertoni, D., Mondini, S., & De Tanti, A. (2023). Long term effects of severe acquired brain injury: A follow-up investigation on the role of cognitive reserve on cognitive outcomes. *Applied Neuropsychology: Adult*, 1-6.

- Baugh, C. M., Stamm, J. M., Riley, D. O., Gavett, B. E., Shenton, M. E., Lin, A., Nowinski, C. J., Cantu, R. C., McKee, A. C., & Stern, R. A. (2012). Chronic traumatic encephalopathy: neurodegeneration following repetitive concussive and subconcussive brain trauma. *Brain imaging and behavior*, 6(2), 244–254.
- Beck, A. T., Epstein, N., Brown, G., & Steer, R. A. (1988). An inventory for measuring clinical anxiety: Psychometric properties. *Journal of Consulting and Clinical Psychology*, 56(6), 893–897.
- Beck, A. T., Steer, R. A., Brown, G. K., (1996). BDI-II, Beck Depression Inventory Manual. *Psychological Corporation*, San Antonio, TX
- Berkman, E. T., Falk, E. B., & Lieberman, M. D. (2011). In the trenches of real-world self-control: neural correlates of breaking the link between craving and smoking. *Psychological Science*, 22(4), 498–506.
- Berkman, E. T., Kahn, L. E., & Merchant, J. S. (2014). Training-induced changes in inhibitory control network activity. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, 34(1), 149–157.
- Berridge, K. C., & Robinson, T. E. (1998). What is the role of dopamine in reward: hedonic impact, reward learning, or incentive salience?. *Brain research. Brain research reviews*, 28(3), 309–369.
- Bianco, F., Lecce, S., & Banerjee, R. (2016). Conversations about mental states and theory of mind development during middle childhood: A training study. *Journal of Experimental Child Psychology*, 149, 41–61.
- Bibby, H., & McDonald, S. (2005). Theory of mind after traumatic brain injury. *Neuropsychologia*, 43(1), 99–114.
- Bigler, E. D., Anderson, C. V., & Blatter, D. D. (2002). Temporal lobe morphology in normal aging and traumatic brain injury. *AJNR. American journal of neuroradiology*, 23(2), 255–266.
- Birmaher, B., Khetarpal, S., Brent, D., Cully, M., Balach, L., Kaufman, J., & Neer, S. M. (1997). The Screen for Child Anxiety Related Emotional Disorders (SCARED): Scale construction and psychometric characteristics. *Journal of the American Academy of Child & Adolescent Psychiatry*, 36(4), 545–553.
- Bivona, U., Costa, A., Contrada, M., Silvestro, D., Azicnuda, E., Aloisi, M., Catania, G., Ciurli, P., Guariglia, C., Caltagirone, C., Formisano, R., & Prigatano, G. P. (2019). Depression, apathy and impaired self-awareness following severe traumatic brain injury: a preliminary investigation. *Brain injury*, 33(9), 1245–1256.
- Blair, R. J., & Cipolotti, L. (2000). Impaired social response reversal. A case of 'acquired sociopathy'. *Brain: a journal of neurology*, 123 (Pt 6), 1122–1141.
- Boumparis, N., Karyotaki, E., Kleiboer, A., Hofmann, S. G., & Cuijpers, P. (2016). The effect of psychotherapeutic interventions on positive and negative affect in depression: A systematic review and meta-analysis. *Journal of affective disorders*, 202, 153–162.
- Brodsky, H., Sachdev, P. S., Withall, A., Altendorf, A., Valenzuela, M. J., & Lorentz, L. (2005). Frequency and clinical, neuropsychological and neuroimaging correlates of apathy following stroke – the Sydney Stroke Study. *Psychological Medicine*, 35(12), 1707–1716.
- Brooks, N., Campsie, L., Symington, C., Beattie, A., & McKinlay, W. (1986). The five year outcome of severe blunt head injury: a relative's view. *Journal of neurology, neurosurgery, and psychiatry*, 49(7), 764–770.
- Broomfield, N. M., Quinn, T. J., Abdul-Rahim, A. H., Walters, M. R., & Evans, J. J. (2014). Depression and anxiety symptoms post-stroke/TIA: Prevalence and associations in cross-sectional data from a regional stroke registry. *BMC Neurology*, 14(1), Article 198.
- Broomfield, N. M., Scoular, A., Welsh, P., Walters, M., & Evans, J. J. (2015). Poststroke anxiety is prevalent at the population level, especially among socially deprived and younger age community stroke survivors. *International Journal of Stroke*, 10(6), 897–902.
- Bryant, R. A., O'Donnell, M. L., Creamer, M., McFarlane, A. C., Clark, C. R., & Silove, D. (2010). The psychiatric sequelae of traumatic injury. *The American journal of psychiatry*, 167(3), 312–320.
- Buijck, B. I., Zuidema, S. U., Spruit-van Eijk, M., Geurts, A. C., & Koopmans, R. T. (2012). Neuropsychiatric symptoms in geriatric patients admitted to skilled nursing facilities in nursing homes for rehabilitation after stroke: a longitudinal multicenter study. *International journal of geriatric psychiatry*, 27(7), 734–741.
- Caciro, L., Ferro, J. M., & Costa, J. (2013). Apathy Secondary to Stroke: A Systematic Review and Meta-Analysis. *Cerebrovascular Diseases*, 35(1), 23–39.
- Calabrò, R. S., Italiano, D., Mili, D., & Bramanti, P. (2012). Levetiracetam-associated loss of libido and anhedonia. *Epilepsy & behavior: E&B*, 24(2), 283–284.
- Camara, E., Rodriguez-Fornells, A., & Münte, T. F. (2009). Functional connectivity of reward processing in the brain. *Frontiers in human neuroscience*, 2, 19.
- Caputi, M., Cugnata, F., & Brombin, C. (2021). Theory of mind and loneliness: Effects of a conversation-based training at school. *International Journal of Psychology*, 56(2), 257–265.
- Cardile, D., Calderone, A., Pagano, M., Cappadona, I., Rifici, C., Quartarone, A., Corallo, F., & Calabrò, R. S. (2024). Coping Strategies in Patients with Acquired Brain Injury: A Scoping Review. *Brain sciences*, 14(8), 784.

- Carver, C. S. (2001). Affect and the Functional Bases of Behavior: On the Dimensional Structure of Affective Experience. *Personality and Social Psychology Review*, 5(4), 345-356.
- Channon, S., Pellieff, A., & Rule, A. (2005). Social cognition after head injury: Sarcasm and theory of mind. *Brain and Language*, 93(2), 123-134.
- Choi, Y., Kim, E. Y., Sun, J., Kim, H. K., Lee, Y. S., Oh, B. M., Park, H. Y., & Leigh, J. H. (2022). Incidence of Depression after Traumatic Brain Injury: A Nationwide Longitudinal Study of 2.2 Million Adults. *Journal of neurotrauma*, 39(5-6), 390-397.
- Choi-Kwon, S., & Kim, J. S. (2022). Anger, a Result and Cause of Stroke: A Narrative Review. *Journal of stroke*, 24(3), 311-322.
- Chun, H.-Y. Y., Whiteley, W. N., Dennis, M. S., Mead, G. E., & Carson, A. J. (2018). Anxiety after stroke: The importance of subtyping. *Stroke*, 49(3), e1-e3.
- Clausi, S., Olivito, G., Lupo, M., Siciliano, L., Bozzali, M., & Leggio, M. (2019). The cerebellar predictions for social interactions: theory of mind abilities in patients with degenerative cerebellar atrophy. *Frontiers in Cellular Neuroscience*, 12, 510.
- Connor, K. M., et al. (2000). Psychometric properties of the Social Phobia Inventory (SPIN): New self-rating scale. *British Journal of Psychiatry*, 176(4), 379-386.
- Cummings, J., Mega, M., Gray, K., Rosenberg-Thompson, S., Carusi, D. A., & Gornbein, J. (1994). The Neuropsychiatric Inventory: Comprehensive assessment of psychopathology in dementia. *Neurology*, 44, 2308-2314.
- Curvis, W., Simpson, J., & Hampson, N. (2016). Social anxiety following traumatic brain injury: An exploration of associated factors. *Neuropsychological Rehabilitation*, 28(4), 527-547.
- Custodio, C. M., & Basford, J. R. (2004). Delayed postanoxic encephalopathy: a case report and literature review. *Archives of Physical Medicine and Rehabilitation*, 85(3), 502-505.
- Damasio, A.R. (1994). The somatic marker hypothesis and the possible functions of the prefrontal cortex. *Philosophical Transactions of the Royal Society of London Biological Sciences*, 351, pp, 1413-1420.
- de Sousa, A., McDonald, S., Rushby, J., Li, S., Dimoska, A., & James, C. (2010). Why don't you feel how I feel? Insight into the absence of empathy after severe traumatic brain injury. *Neuropsychologia*, 48(12), 3585-3595.
- Dehbozorgi, M., Maghsoudi, M. R., Mohammadi, I., et al. (2024). Incidence of anxiety after traumatic brain injury: A systematic review and meta-analysis. *BMC Neurology*, 24, 293.
- Der-Avakian, A., & Markou, A. (2012). The neurobiology of anhedonia and other reward-related deficits. *Trends in neurosciences*, 35(1), 68-77.
- Diamond, P. R., Dysch, L., & Daniels, J. (2023). Health anxiety in stroke survivors: a cross-sectional study on the prevalence of health anxiety in stroke survivors and its impact on quality of life. *Disability and rehabilitation*, 45(1), 27-33.
- Douglas, J. M., & Spellacy, F. J. (1996). Indicators of long-term family functioning following severe traumatic brain injury in adults. *Brain injury*, 10(11), 819-839.
- Dunn, B. D., German, R. E., Khazanov, G. K., Xu, C., Hollon, S. D., & DeRubeis, R. J. (2019). Changes in Positive and Negative Affect During Pharmacological Treatment and Cognitive Therapy for Major Depressive Disorder: A Secondary Analysis of Two Randomized Controlled Trials. *Clinical Psychological Science*, 8, 36 - 51
- Eslinger, P. J., & Damasio, A. R. (1985). Severe disturbance of higher cognition after bilateral frontal lobe ablation: patient EVR. *Neurology*, 35(12), 1731-1741.
- Fedoroff, J. P., Starkstein, S. E., Forrester, A. W., Geisler, F. H., Jorge, R. E., Arndt, S. V., & Robinson, R. G. (1992). Depression in patients with acute traumatic brain injury. *The American journal of psychiatry*, 149(7), 918-923.
- Feng, C., Fang, M., & Liu, X. Y. (2014). The neurobiological pathogenesis of poststroke depression. *TheScientificWorldJournal*, 2014, 521349.
- Frank, D., Gruenbaum, B. F., Zlotnik, A., Semyonov, M., Frenkel, A., & Boyko, M. (2022). Pathophysiology and Current Drug Treatments for Post-Stroke Depression: A Review. *International journal of molecular sciences*, 23(23), 15114.
- Frith, U., & Frith, C. D. (2003). Development and neurophysiology of mentalizing. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 358(1431), 459-473.
- Gainotti, G., Antonucci, G., Marra, C., & Paolucci, S. (2001). Relation between depression after stroke, antidepressant therapy, and functional recovery. *Journal of neurology, neurosurgery, and psychiatry*, 71(2), 258-261.
- Gard, D. E., Kring, A. M., Gard, M. G., Horan, W. P., & Green, M. F. (2007). Anhedonia in schizophrenia: distinctions between anticipatory and consummatory pleasure. *Schizophrenia research*, 93(1-3), 253-260.
- Gard, D. E., Gard, M. G., Kring, A. M., & John, O. P. (2006). Anticipatory and consummatory components of the experience of pleasure: A scale development study. *Journal of Research in Personality*, 40(6), 1086-1102.

- Ghika-Schmid, F., van Melle, G., Guex, P., & Bogousslavsky, J. (1999). Subjective experience and behavior in acute stroke: the Lausanne Emotion in Acute Stroke Study. *Neurology*, *52*(1), 22–28.
- Giles, G. M., & Mohr, J. D. (2007). Overview and inter-rater reliability of an incident-based rating scale for aggressive behavior following traumatic brain injury: The Overt Aggression Scale-Modified for Neurorehabilitation-Extended (OAS-MNR-E). *Brain injury*, *21*(5), 505–511.
- Goldman, L., Siddiqui, E. M., Khan, A., Jahan, S., Rehman, M. U., Mehan, S., Sharma, R., Budkin, S., Kumar, S. N., Sahu, A., Kumar, M., & Vaibhav, K. (2022). Understanding Acquired Brain Injury: A Review. *Biomedicines*, *10*(9), 2167.
- Green, S. L., Gignac, G. E., Watson, P. A., Brosnan, N., Becerra, R., Pestell, C., & Weinborn, M. (2022). Apathy and Depression as Predictors of Activities of Daily Living Following Stroke and Traumatic Brain Injuries in Adults: A Meta-Analysis. *Neuropsychology review*, *32*(1), 51–69.
- Greenop, K. R., Almeida, O. P., Hankey, G. J., van Bockxmeer, F., & Lautenschlager, N. T. (2009). Premorbid personality traits are associated with post-stroke behavioral and psychological symptoms: a three-month follow-up study in Perth, Western Australia. *International psychogeriatrics*, *21*(6), 1063–1071.
- Hamilton, J., Radlak, B., Morris, P. G., & Phillips, L. H. (2017). Theory of mind and executive functioning following stroke. *Archives of Clinical Neuropsychology*, *32*(5), 507–518.
- Hart, T., Fann, J. R., Brockway, J. A., Maiuro, R. D. (2017). Treatment of anger and aggression following acquired brain injury. In: Sturmeijer P, Ed., *The Wiley Handbook of Violence and Aggression*. New York: John Wiley & Sons, 2017, chapter 80.
- Hartman-Macir, A., Soroker, N., Ring, H., Avni, N., & Katz, N. (2007). Activities, participation and satisfaction one-year post stroke. *Disability and rehabilitation*, *29*(7), 559–566.
- Hathaway, S. R., & McKinley, J. C. (1940). A multiphasic personality schedule (Minnesota): I. Construction of the schedule. *The Journal of Psychology*, *10*(2), 249–254.
- He, N., Rolls, E.T., Zhao, W. *et al.* Predicting human inhibitory control from brain structural MRI. *Brain Imaging and Behavior*, *14*, 2148–2158 (2020).
- Henry, J. D., Phillips, L. H., Crawford, J. R., Ietswaart, M., & Summers, F. (2006). Theory of mind following traumatic brain injury: The role of emotion recognition and executive dysfunction. *Neuropsychologia*, *44*(10), 1623–1628.
- Hinkeldey, N. S., & Corrigan, J. D. (1990). The structure of head-injured patients' neurobehavioural complaints: a preliminary study. *Brain injury*, *4*(2), 115–133.
- Ho, N., & Sommers, M. (2013). Anhedonia: a concept analysis. *Archives of psychiatric nursing*, *27*(3), 121–129.
- Howlett, J. R., Nelson, L. D., & Stein, M. B. (2022). Mental Health Consequences of Traumatic Brain Injury. *Biological psychiatry*, *91*(5), 413–420.
- Hudgens, S., Floden, L., Blackowicz, M., Jamieson, C., Popova, V., Fedgchin, M., Drevets, W. C., Cooper, K., Lane, R., & Singh, J. (2021). Meaningful Change in Depression Symptoms Assessed with the Patient Health Questionnaire (PHQ-9) and Montgomery-Åsberg Depression Rating Scale (MADRS) Among Patients with Treatment Resistant Depression in Two, Randomized, Double-blind, Active-controlled Trials of Esketamine Nasal Spray Combined with a New Oral Antidepressant. *Journal of affective disorders*, *281*, 767–775.
- Ito, D., Mori, N., Shimizu, A., Narita, A., Sakata, S., Honaga, K., Kondo, K., & Otaka, Y. (2023). Presence and Characteristics of Behavioral and Psychological Symptoms in Subacute Stroke Patients with Cognitive Impairment. *Behavioural neurology*, *2023*, 6636217.
- Iverson, G. L., & Gardner, A. J. (2021). Symptoms of traumatic encephalopathy syndrome are common in the US general population. *Brain communications*, *3*(1), feab001.
- Jenkins, L. M., Heywood, A., Gupta, S., Kouchakidivkolaei, M., Sridhar, J., Rogalski, E., Weintraub, S., Popuri, K., Rosen, H., Wang, L., & Frontotemporal Lobar Degeneration Neuroimaging Initiative (2024). Disinhibition in dementia related to reduced morphometric similarity of cognitive control network. *Brain communications*, *6*(2).
- Jorge, R. E., Robinson, R. G., Arndt, S. V., Forrester, A. W., Geisler, F., & Starkstein, S. E. (1993). Comparison between acute- and delayed-onset depression following traumatic brain injury. *The Journal of neuropsychiatry and clinical neurosciences*, *5*(1), 43–49.
- Jorge, R. E., & Starkstein, S. E. (2005). Pathophysiologic aspects of major depression following traumatic brain injury. *The Journal of head trauma rehabilitation*, *20*(6), 475–487.
- Jorge, R. E., Starkstein, S. E., & Robinson, R. G. (2010). Apathy following Stroke. *The Canadian Journal of Psychiatry*, *55*(6), 350–354.
- Juengst, S. B., Kumar, R. G., Arenth, P. M., & Wagner, A. K. (2014). Exploratory associations with tumor necrosis factor- α , disinhibition and suicidal endorsement after traumatic brain injury. *Brain, behavior, and immunity*, *41*, 134–143.
- Kant, R., Duffy, J. D., & Pivovarnik, A. (1998). Prevalence of apathy following head injury. *Brain Injury*, *12*(1), 87–92.

- Kashdan, T. B., Weeks, J. W., & Savostyanova, A. A. (2011). Whether, how, and when social anxiety shapes positive experiences and events: a self-regulatory framework and treatment implications. *Clinical psychology review, 31*(5), 786–799.
- Keedwell, P. A., Andrew, C., Williams, S. C., Brammer, M. J., & Phillips, M. L. (2005). The neural correlates of anhedonia in major depressive disorder. *Biological psychiatry, 58*(11), 843–853.
- Kelly, G., Todd, J., Simpson, G., Kremer, P., & Martin, C. (2006). The Overt Behaviour Scale (OBS): a tool for measuring challenging behaviours following ABI in community settings. *Brain injury, 20*(3), 307–319.
- Kelly, G., Brown, S., Todd, J., & Kremer, P. (2008). Challenging behaviour profiles of people with acquired brain injury living in community settings. *Brain injury, 22*(6), 457–470.
- Kempton, M. J., Salvador, Z., Munafò, M. R., Geddes, J. R., Simmons, A., Frangou, S., & Williams, S. C. (2011). Structural neuroimaging studies in major depressive disorder. Meta-analysis and comparison with bipolar disorder. *Archives of general psychiatry, 68*(7), 675–690.
- Kendall, A. D., Zinbarg, R. E., Mineka, S., Bobova, L., Prenoveau, J. M., Revelle, W., & Craske, M. G. (2015). Prospective associations of low positive emotionality with first onsets of depressive and anxiety disorders: Results from a 10-wave latent trait-state modeling study. *Journal of abnormal psychology, 124*(4), 933–943.
- Khazanov, G. K., & Ruscio, A. M. (2016). Is low positive emotionality a specific risk factor for depression? A meta-analysis of longitudinal studies. *Psychological bulletin, 142*(9), 991–1015.
- Khazanov, G. K., Ruscio, A. M., & Forbes, C. N. (2020). The Positive Valence Systems Scale: Development and Validation. *Assessment, 27*(5), 1045–1069.
- Kim E. (2002). Agitation, aggression, and disinhibition syndromes after traumatic brain injury. *NeuroRehabilitation, 17*(4), 297–310.
- Kim J. S. (2016). Post-stroke Mood and Emotional Disturbances: Pharmacological Therapy Based on Mechanisms. *Journal of stroke, 18*(3), 244–255.
- Kim, J. S., Choi, S., Kwon, S. U., & Seo, Y. S. (2002). Inability to control anger or aggression after stroke. *Neurology, 58*(7), 1106–1108.
- Knutson, K. M., Dal Monte, O., Schintu, S., Wassermann, E. M., Raymont, V., Grafman, J., & Krueger, F. (2015). Areas of Brain Damage Underlying Increased Reports of Behavioral Disinhibition. *The Journal of neuropsychiatry and clinical neurosciences, 27*(3), 193–198.
- Kootker, J. A., van Mierlo, M. L., Hendriks, J. C., et al. (2016). Risk factors for symptoms of depression and anxiety one year post-stroke: A longitudinal study. *Archives of Physical Medicine and Rehabilitation, 97*(6), 919–928.
- Korkiakangas, T., Dindar, K., Laitila, A., & Kärnä, E. (2016). The Sally–Anne test: an interactional analysis of a dyadic assessment. *International journal of language & communication disorders, 51*(6), 685–702.
- Koskinen S. (1998). Quality of life 10 years after a very severe traumatic brain injury (TBI): the perspective of the injured and the closest relative. *Brain injury, 12*(8), 631–648.
- Lamontagne, G., Belleville, G., Beaulieu-Bonneau, S., Souesme, G., Savard, J., Sirois, M.-J., Giguère, M., Tessier, D., Le Sage, N., & Ouellet, M.-C. (2022). Anxiety symptoms and disorders in the first year after sustaining mild traumatic brain injury. *Rehabilitation Psychology, 67*(1), 90–99.
- Lau, C. G., Tang, W. K., Liu, X. X., Liang, H. J., Liang, Y., Wong, A., Mok, V., Ungvari, G. S., Wong, K. S., Kim, J. S., & Paradiso, S. (2017). Poststroke agitation and aggression and social quality of life: a case control study. *Topics in stroke rehabilitation, 24*(2), 126–133.
- Lecce, S., Bianco, F., Devine, R. T., Hughes, C., & Banerjee, R. (2014). Promoting theory of mind during middle childhood: A training program. *Journal of experimental child psychology, 126*, 52–67.
- Levy, R., & Dubois, B. (2005). Apathy and the functional anatomy of the prefrontal Cortex–Basal ganglia circuits. *Cerebral Cortex, 16*(7), 916–928.
- Lewis, J. D., Krueger, F., Raymont, V., Solomon, J., Knutson, K. M., Barbey, A. K., Poore, J. C., Wassermann, E. M., & Grafman, J. (2015). Anhedonia in combat veterans with penetrating head injury. *Brain Imaging and Behavior, 9*(3), 456–460.
- Lezak, M. D., Howieson, D. B., Loring, D. W., Hannay, H. J., & Fischer, J. S. (2004). *Neuropsychological assessment* (4th ed.). Oxford University Press.
- Liu, F., Yang, Y., Wang, S., et al. (2021). Impact of sleep duration on depression and anxiety after acute ischemic stroke. *Frontiers in Neurology, 12*, 630–638.
- Llerena, K., Park, S. G., McCarthy, J. M., Couture, S. M., Bennett, M. E., & Blanchard, J. J. (2013). The Motivation and Pleasure Scale-Self-Report (MAP-SR): reliability and validity of a self-report measure of negative symptoms. *Comprehensive psychiatry, 54*(5), 568–574.
- Mallorquí, A., Martínez-Zamora, M. A., & Carmona, F. (2022). Anhedonia in endometriosis: An unexplored symptom. *Frontiers in psychology, 13*, 935349.

- Marcheschi, E., Von Koch, L., Pessah-Rasmussen, H., & Elf, M. (2018). Home setting after stroke, facilitators and barriers: A systematic literature review. *Health & social care in the community*, 26(4), e451–e459.
- Marin, R. S. (1991). Apathy: a neuropsychiatric syndrome. *The Journal of neuropsychiatry and clinical neurosciences*, 3(3), 243–254.
- Marin, R. S., Biedrzycki, R. C., & Firinciogullari, S. (1991). Reliability and validity of the Apathy Evaluation Scale. *Psychiatry research*, 38(2), 143–162.
- Marks, M. R., Dux, M. C., Rao, V., & Albrecht, J. S. (2021). Treatment patterns of anxiety and posttraumatic stress disorder following traumatic brain injury. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 34(3).
- Martin, I., & McDonald, S. (2005). Evaluating the causes of impaired irony comprehension following traumatic brain injury. *Aphasiology*, 19(8), 712–730.
- Mata-Bermudez, A., Trejo-Chávez, R., Martínez-Vargas, M., Pérez-Arredondo, A., Martínez-Cardenas, M. L. Á., Diaz-Ruiz, A., Rios, C., & Navarro, L. (2024). Dysregulation of the dopaminergic system secondary to traumatic brain injury: Implications for mood and anxiety disorders. *Frontiers in Neuroscience*, 18, 1447688.
- Mayberg, H. S., Liotti, M., Brannan, S. K., McGinnis, S., Mahurin, R. K., Jerabek, P. A., Silva, J. A., Tekell, J. L., Martin, C. C., Lancaster, J. L., & Fox, P. T. (1999). Reciprocal limbic-cortical function and negative mood: converging PET findings in depression and normal sadness. *The American journal of psychiatry*, 156(5), 675–682.
- McDonald, S., & Flanagan, S. (2004). Social perception deficits after traumatic brain injury: interaction between emotion recognition, mentalizing ability, and social communication. *Neuropsychology*, 18(3), 572.
- McDonald, S. (2013). Impairments in social cognition following severe traumatic brain injury. *Journal of the International Neuropsychological Society*, 19(3), 231–246.
- McIntosh, T. K., Saatman, K. E., Raghupathi, R., Graham, D. I., Smith, D. H., Lee, V. M., & Trojanowski, J. Q. (1998). The Dorothy Russell Memorial Lecture. The molecular and cellular sequelae of experimental traumatic brain injury: pathogenetic mechanisms. *Neuropathology and applied neurobiology*, 24(4), 251–267.
- Menon, D. K., Schwab, K., Wright, D. W., & Maas, A. I. (2010). Position Statement: Definition of Traumatic Brain Injury. *Archives of Physical Medicine and Rehabilitation*, 91(11), 1637–1640.
- Mok, V. C., Wong, A., Wong, K., Chu, W. C., Xiong, Y., Chan, A. Y., Kwok, T. C., Hu, X., Lee, W. K., Tang, W. K., Wong, K. S., & Wong, S. (2010). Executive dysfunction and left frontal white matter hyperintensities are correlated with neuropsychiatric symptoms in stroke patients with confluent white matter hyperintensities. *Dementia and geriatric cognitive disorders*, 30(3), 254–260.
- Montenigro, P. H., Baugh, C. M., Daneshvar, D. H., Mez, J., Budson, A. E., Au, R., Katz, D. I., Cantu, R. C., & Stern, R. A. (2014). Clinical subtypes of chronic traumatic encephalopathy: literature review and proposed research diagnostic criteria for traumatic encephalopathy syndrome. *Alzheimer's research & therapy*, 6(5), 68.
- Montgomery, S. A., & Asberg, M. (1979). A new depression scale designed to be sensitive to change. *The British journal of psychiatry: the journal of mental science*, 134, 382–389.
- Morey, L. C., & Ambwani, S. (2008). The Personality Assessment Inventory. In G. J. Boyle, G. Matthews, & D. H. Saklofske (Eds.), *The SAGE handbook of personality theory and assessment, Vol. 2. Personality measurement and testing* (pp. 626–645). Sage Publications, Inc.
- Morris, B. H., Bylsma, L. M., & Rottenberg, J. (2009). Does emotion predict the course of major depressive disorder? A review of prospective studies. *The British journal of clinical psychology*, 48(Pt 3), 255–273.
- Nawijn, L., van Zuiden, M., Frijling, J. L., Koch, S. B., Veltman, D. J., & Olf, M. (2015). Reward functioning in PTSD: a systematic review exploring the mechanisms underlying anhedonia. *Neuroscience and biobehavioral reviews*, 51, 189–204.
- Neumann, D., Malec, J. F., & Hammond, F. M. (2015). The association of negative attributions with irritation and anger after brain injury. *Rehabilitation psychology*, 60(2), 155–161.
- Oliva, G., Masina, F., Hosseinkhani, N., Montemurro, S., & Arcara, G. (2024). Cognitive reserve in the recovery and rehabilitation of stroke and traumatic brain injury: A systematic review. *The Clinical neuropsychologist*, 1–37. Advance online publication.
- Omalu, B., Bailes, J., Hamilton, R. L., Kamboh, M. I., Hammers, J., Case, M., & Fitzsimmons, R. (2011). Emerging histomorphologic phenotypes of chronic traumatic encephalopathy in American athletes. *Neurosurgery*, 69(1), 173–183.
- Osborn, A. J., Mathias, J. L., & Fairweather-Schmidt, A. K. (2016). Prevalence of anxiety following adult traumatic brain injury: A meta-analysis comparing measures, samples and postinjury intervals. *Neuropsychology*, 30(2), 247–261.
- Osborne-Crowley, K., & McDonald, S. (2018). A review of social disinhibition after traumatic brain injury. *Journal of neuropsychology*, 12(2), 176–199.

- Paolucci, S., Iosa, M., Coiro, P., Venturiero, V., Savo, A., De Angelis, D., & Morone, G. (2019). Post-stroke Depression Increases Disability More Than 15% in Ischemic Stroke Survivors: A Case-Control Study. *Frontiers in neurology, 10*, 926.
- Paradiso, S., Robinson, R. G., & Arndt, S. (1996). Self-reported aggressive behavior in patients with stroke. *The Journal of nervous and mental disease, 184*(12), 746–753.
- Patel, M. D., Coshall, C., Rudd, A. G., & Wolfe, C. D. (2002). Cognitive impairment after stroke: clinical determinants and its associations with long-term stroke outcomes. *Journal of the American Geriatrics Society, 50*(4), 700–706.
- Paul, R. H., Brickman, A. M., Navia, B., Hinkin, C., Malloy, P. F., Jefferson, A. L., Cohen, R. A., Tate, D. F., & Flanigan, T. P. (2005). Apathy is Associated with Volume of the Nucleus Accumbens in Patients Infected With HIV. *Journal of Neuropsychiatry, 17*(2), 167–171.
- Pelizza, L., & Ferrari, A. (2009). Anhedonia in schizophrenia and major depression: state or trait?. *Annals of general psychiatry, 8*, 22.
- Petrova, E. A., Savina, M. A., Kontsevoi, V. A., & Skvortsova, V. I. (2012). Clinical characteristics of post-stroke anxiety disorders. *Klinika Nervnykh I Psikhicheskikh Zabolevaniï, 9*, 12–16.
- Piamarta, F., Iurlaro, S., Isella, V., Atzeni, L., Grimaldi, M., Russo, A.C., Forapani, E., & Appollonio, I. (2004). Unconventional affective symptoms and executive functions after stroke in the elderly. *Archives of gerontology and geriatrics. Supplement, 9*, 315-23.
- Preda, A. (2014). Brain Imaging Correlates of Anhedonia. In: Ritsner, M. (eds) *Anhedonia: A Comprehensive Handbook Volume I*. Springer, Dordrecht.
- Radakovic, R., Stephenson, L., Colville, S., Swingler, R., Chandran, S., & Abrahams, S. (2016). Multidimensional apathy in ALS: validation of the Dimensional Apathy Scale. *Journal of neurology, neurosurgery, and psychiatry, 87*(6), 663–669.
- Rafsten, L., Danielsson, A., & Sunnerhagen, K. S. (2018). Anxiety after stroke: A systematic review and meta-analysis. *Journal of Rehabilitation Medicine, 50*(9), 769–778.
- Rao, V., Rosenberg, P., Bertrand, M., Salehina, S., Spiro, J., Vaishnavi, S., Rastogi, P., Noll, K., Schretlen, D. J., Brandt, J., Cornwell, E., Makley, M., & Miles, Q. S. (2009). Aggression after traumatic brain injury: prevalence and correlates. *The Journal of neuropsychiatry and clinical neurosciences, 21*(4), 420–429.
- Rao, V., Spiro, J. R., Schretlen, D. J., & Cascella, N. G. (2007). Apathy syndrome after traumatic brain injury compared with deficits in schizophrenia. *Psychosomatics, 48*(3), 217–222.
- Rao, V., & Lyketsos, C. (2000). Neuropsychiatric sequelae of traumatic brain injury. *Psychosomatics, 41*(2), 95–103.
- Raymont, V., Salazar, A. M., Krueger, F., & Grafman, J. (2011). "Studying injured minds" - the Vietnam head injury study and 40 years of brain injury research. *Frontiers in neurology, 2*, 15.
- Ren, J., Friedmann, D., Xiong, J., Liu, C. D., Ferguson, B. R., Weerakkody, T., DeLoach, K. E., Ran, C., Pun, A., Sun, Y., Weissbourd, B., Neve, R. L., Huguenard, J., Horowitz, M. A., & Luo, L. (2018). Anatomically Defined and Functionally Distinct Dorsal Raphe Serotonin Sub-systems. *Cell, 175*(2), 472–487.e20.
- Rizvi, S. J., Quilty, L. C., Sproule, B. A., Cyriac, A., Michael Bagby, R., & Kennedy, S. H. (2015). Development and validation of the Dimensional Anhedonia Rating Scale (DARS) in a community sample and individuals with major depression. *Psychiatry research, 229*(1-2), 109–119.
- Roth, R. M., Flashman, L. A., Saykin, A. J., McAllister, T. W., & Vidaver, R. (2004). Apathy in schizophrenia: reduced frontal lobe volume and neuropsychological deficits. *American Journal of Psychiatry, 161*(1), 157–159.
- Roy, D., Vaishnavi, S., Han, D., & Rao, V. (2017). Correlates and Prevalence of Aggression at Six Months and One Year After First-Time Traumatic Brain Injury. *The Journal of neuropsychiatry and clinical neurosciences, 29*(4), 334–342.
- Sacher, J., Neumann, J., Fünfstück, T., Soliman, A., Villringer, A., & Schroeter, M. L. (2012). Mapping the depressed brain: a meta-analysis of structural and functional alterations in major depressive disorder. *Journal of affective disorders, 140*(2), 142–148.
- Sandman, C. F., & Craske, M. G. (2022). Psychological Treatments for Anhedonia. *Current topics in behavioral neurosciences, 58*, 491–513.
- Santos, C. O., Caeiro, L., Ferro, J. M., Albuquerque, R., & Luísa Figueira, M. (2006). Anger, hostility and aggression in the first days of acute stroke. *European journal of neurology, 13*(4), 351–358.
- Saxe, R., Carey, S., & Kanwisher, N. (2004). Understanding other minds: linking developmental psychology and functional neuroimaging. *Annual Review of Psychology, 55*(1), 87-124.
- Schultz, W. (2002). Getting formal with dopamine and reward. *Neuron, 36*(2), 241–263.
- Segura, E., Vilà-Balló, A., Mallorquí, A., Porto, M. F., Duarte, E., Grau-Sánchez, J., & Rodríguez-Fornells, A. (2024). The presence of anhedonia in individuals with subacute and chronic stroke: an exploratory cohort study. *Frontiers in aging neuroscience, 16*, 1253028.

- Shamay-Tsoory, S. G., Tomer, R., Goldsher, D., Berger, B. D., & Aharon-Peretz, J. (2004). Impairment in cognitive and affective empathy in patients with brain lesions: anatomical and cognitive correlates. *Journal of Clinical and Experimental Neuropsychology*, *26*(8), 1113-1127.
- Sibon, I., Lassalle-Lagadec, S., Renou, P., & Swendsen, J. (2012). Evolution of depression symptoms following stroke: a prospective study using computerized ambulatory monitoring. *Cerebrovascular diseases (Basel, Switzerland)*, *33*(3), 280-285.
- Simpson, G., Blaszczyński, A., & Hodgkinson, A. (1999). Sex offending as a psychosocial sequela of traumatic brain injury. *The Journal of head trauma rehabilitation*, *14*(6), 567-580.
- Söderstrand, P., & Almkvist, O. (2012). Psychometric data on the Eyes Test, the Faux Pas Test, and the Dewey Social Stories Test in a population-based Swedish adult sample. *Nordic Psychology*, *64*(1), 30-43.
- Soo, C., Kiernan, M., & Anderson, V. (2019). Trait mindfulness as a mediator of anxiety and psychosocial functioning in young people with acquired brain injury. *Developmental Neurorehabilitation*, *23*(4), 231-239.
- Spatt, J., Zebenholzer, K., & Oder, W. (1997). Psychosocial long-term outcome of severe head injury as perceived by patients, relatives, and professionals. *Acta Neurologica Scandinavica*, *95*(3), 173-179.
- Spielberger, C. D. (1989). *State-Trait Anxiety Inventory: Bibliography* (2nd ed.). Consulting Psychologists Press.
- Starkstein, S. E., Fedoroff, J. P., Price, T. R., Leiguarda, R., & Robinson, R. G. (1993). Apathy following cerebrovascular lesions. *Stroke*, *24*(11), 1625-1630.
- Starkstein, S. E., Mayberg, H. S., Preziosi, T. J., Andrezejewski, P., Leiguarda, R., & Robinson, R. G. (1992). Reliability, validity, and clinical correlates of apathy in Parkinson's disease. *The Journal of neuropsychiatry and clinical neurosciences*, *4*(2), 134-139.
- Stern, Y., Albert, M., Barnes, C. A., Cabeza, R., Pascual-Leone, A., & Rapp, P. R. (2023). A framework for concepts of reserve and resilience in aging. *Neurobiology of aging*, *124*, 100-103.
- Stone, V. E., Baron-Cohen, S., & Knight, R. T. (1998). Frontal lobe contributions to theory of mind. *Journal of Cognitive Neuroscience*, *10*(5), 640-656.
- Tang, E. Y. H., Price, C., Stephan, B. C. M., Robinson, L., & Exley, C. (2020). Impact of Memory Problems Post-stroke on Patients and Their Family Carers: A Qualitative Study. *Frontiers in medicine*, *7*, 267.
- Tateno, A., Jorge, R. E., & Robinson, R. G. (2003). Clinical correlates of aggressive behavior after traumatic brain injury. *The Journal of neuropsychiatry and clinical neurosciences*, *15*(2), 155-160.
- Tay, J., Morris, R. G., & Markus, H. S. (2021). Apathy after stroke: Diagnosis, mechanisms, consequences, and treatment. *International journal of stroke: official journal of the International Stroke Society*, *16*(5), 510-518.
- Terroni, L., Amaro, E., Jr, Iosifescu, D. V., Mattos, P., Yamamoto, F. I., Tinone, G., Conforto, A. B., Sobreiro, M. F., Guajardo, V. D., De Lucia, M. C., Moreira, A. C., Scaff, M., Leite, C. C., & Fraguas, R. (2015). The association of post-stroke anhedonia with salivary cortisol levels and stroke lesion in hippocampal/parahippocampal region. *Neuropsychiatric disease and treatment*, *11*, 233-242.
- Rømer Thomsen, K., Whybrow, P. C., & Kringelbach, M. L. (2015). Reconceptualizing anhedonia: novel perspectives on balancing the pleasure networks in the human brain. *Frontiers in behavioral neuroscience*, *9*, 49.
- Treadway, M. T., & Zald, D. H. (2011). Reconsidering anhedonia in depression: lessons from translational neuroscience. *Neuroscience and biobehavioral reviews*, *35*(3), 537-555.
- Tremblay, L. K., Naranjo, C. A., Cardenas, L., Herrmann, N., & Busto, U. E. (2002). Probing brain reward system function in major depressive disorder: altered response to dextroamphetamine. *Archives of general psychiatry*, *59*(5), 409-416.
- Tremblay, L. K., Naranjo, C. A., Graham, S. J., Herrmann, N., Mayberg, H. S., Hevenor, S., et al. (2005). Functional neuroanatomical substrates of altered reward processing in major depressive disorder revealed by a dopaminergic probe. *Archives of General Psychiatry*, *62*(11), 1228-1236.
- Tso, I. F., Grove, T. B., & Taylor, S. F. (2014). Differential hedonic experience and behavioral activation in schizophrenia and bipolar disorder. *Psychiatry Research*, *219*(3), 470-476.
- Turkstra, L. S., Dixon, T. M., & Baker, K. K. (2004). Theory of mind and social beliefs in adolescents with traumatic brain injury. *NeuroRehabilitation*, *19*(3), 245-256.
- van Almenkerk, S., Depla, M. F., Smalbrugge, M., Eefsting, J. A., & Hertogh, C. M. (2012). Institutionalized stroke patients: status of functioning of an under researched population. *Journal of the American Medical Directors Association*, *13*(7), 634-639.
- Van Zomeran, A. H., & van den Burg, W. (1985). Residual complaints of patients two years after severe head injury. *Journal of neurology, neurosurgery, and psychiatry*, *48*(1), 21-28.
- Verberne, D. P. J., Spauwen, P. J. J., & van Heugten, C. M. (2019). Psychological interventions for treating neuropsychiatric consequences of acquired brain injury: A systematic review. *Neuropsychological rehabilitation*, *29*(10), 1509-1542.

- Vinckier, F., Gourion, D., & Mouchabac, S. (2017). Anhedonia predicts poor psychosocial functioning: Results from a large cohort of patients treated for major depressive disorder by general practitioners. *European psychiatry: the journal of the Association of European Psychiatrists*, *44*, 1–8.
- Visscher, A. J., van Meijel, B., Stolker, J. J., Wiersma, J., & Nijman, H. (2011). Aggressive behaviour of inpatients with acquired brain injury. *Journal of clinical nursing*, *20*(23-24), 3414–3422.
- Warriner, E. M., Rourke, B. P., Velikonja, D., & Metham, L. (2003). Subtypes of emotional and behavioural sequelae in patients with traumatic brain injury. *Journal of clinical and experimental neuropsychology*, *25*(7), 904–917.
- Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures of positive and negative affect: the PANAS scales. *Journal of personality and social psychology*, *54*(6), 1063–1070.
- Watson, D., & Naragon-Gainey, K. (2010). On the specificity of positive emotional dysfunction in psychopathology: evidence from the mood and anxiety disorders and schizophrenia/schizotypy. *Clinical psychology review*, *30*(7), 839–848.
- Whyte, E. M., & Mulsant, B. H. (2002). Post stroke depression: epidemiology, pathophysiology, and biological treatment. *Biological psychiatry*, *52*(3), 253–264.
- Wichers, M., Peeters, F., Geschwind, N., Jacobs, N., Simons, C. J., Derom, C., Thiery, E., Delespaul, P. H., & van Os, J. (2010). Unveiling patterns of affective responses in daily life may improve outcome prediction in depression: a momentary assessment study. *Journal of affective disorders*, *124*(1-2), 191–195.
- Winer, E. S., Nadorff, M. R., Ellis, T. E., Allen, J. G., Herrera, S., & Salem, T. (2014). Anhedonia predicts suicidal ideation in a large psychiatric inpatient sample. *Psychiatry research*, *218*(1-2), 124–128.
- Winer, E. S., Veilleux, J. C., & Ginger, E. J. (2014). Development and validation of the Specific Loss of Interest and Pleasure Scale (SLIPS). *Journal of affective disorders*, *152-154*, 193–201.
- Wise R. A. (2008). Dopamine and reward: the anhedonia hypothesis 30 years on. *Neurotoxicity research*, *14*(2-3), 169–183.
- Wong, A., Cheng, S. T., Lo, E. S., Kwan, P. W., Law, L. S., Chan, A. Y., Wong, L. K., & Mok, V. (2014). Validity and reliability of the neuropsychiatric inventory questionnaire version in patients with stroke or transient ischemic attack having cognitive impairment. *Journal of geriatric psychiatry and neurology*, *27*(4), 247–252.
- World Health Organization. (2004). *International statistical classification of diseases and related health problems (10th rev., 2nd ed.)*. World Health Organization.
- Yeh, S.-C. J., & Lo, S. K. (2004). Living alone, social support, and feeling lonely among the elderly. *Social Behavior and Personality: An International Journal*, *32*(2), 129–138.
- Yeh, Z. T., & Tsai, C. F. (2014). Impairment on theory of mind and empathy in patients with stroke. *Psychiatry and Clinical Neurosciences*, *68*(8), 612–620.
- Zasler, N. D., Katz, D. I., Zafonte, R. D., Arciniegas, D. B., Bullock, M. R., & Kreutzer, J. S. (Eds.). (2012). *Brain Injury Medicine, 2nd Edition: Principles and Practice*. New York, NY, US: Demos Medical.
- Zigmond, A. S., & Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica*, *67*(6), 361–370.

Gli effetti di un intervento di psicologia positiva sul benessere psicologico degli adolescenti: uno studio preliminare

Effects of positive psychology intervention on the psychological well-being of adolescents: a preliminary study

Mariachiara Gaita^{1*}, Mariamichela Aquino^{1*}, Alfredo Di Dio², Maria Cropano¹, Sabatino Galietta¹, Antonella Fiorentino¹, Simona Raimo³, Florindo d'Onofrio^{2,4}

1. Dipartimento di Psicologia, Università degli Studi della Campania "Luigi Vanvitelli", Caserta, Italia

2. Rotary Club Avellino, Avellino, Italia

3. Dipartimento di Scienze Mediche e Chirurgiche, Università "Magna Graecia" di Catanzaro, Catanzaro, Italia

4. U.O.S.D. Stroke Unit. AORN "S. G. Moscati", Avellino, Italia

*Mariachiara Gaita e Mariamichela Aquino sono co-primi autori

Abstract

L'adolescenza è una fase critica dello sviluppo caratterizzata da cambiamenti significativi che possono influenzare il benessere psicologico e la salute mentale. In questo contesto, il presente studio ha esaminato l'efficacia di un intervento di Psicologia Positiva, in accordo alle linee guida di Seligman (2002), sui fattori protettivi del disagio psicologico (intelligenza emotiva, resilienza, competenze interpersonali) in un campione di 45 adolescenti di età compresa tra i 13 e i 16 anni. Tutti i partecipanti hanno completato una batteria di questionari standardizzati prima (T0) e dopo (T1) l'intervento di Psicologia Positiva. In generale, i risultati mostrano miglioramenti significativi nei punteggi ottenuti ai questionari valutanti le abilità relate all'intelligenza emotiva e le competenze in ambito interpersonale, sottolineando come la Psicologia Positiva sia in grado di rendere gli adolescenti maggiormente capaci di riconoscere, comprendere e gestire le proprie ed altrui emozioni, con riduzioni significative del disagio emotivo provato nelle situazioni sociali. Gli interventi di Psicologia Positiva in adolescenza contribuiscono a rafforzare aspetti psicologici protettivi, risultando così utili nella prevenzione del disagio psicologico.

Parole chiave

Psicologia positiva; Benessere psicologico; Adolescenza; Qualità di vita

Autore responsabile per la corrispondenza: Mariachiara Gaita, Dipartimento di Psicologia, Università degli Studi della Campania "Luigi Vanvitelli", Caserta, Italia, e-mail: mariachiara.gaita@unicampania.it

Abstract

Adolescence is a critical developmental stage characterized by significant changes that can influence psychological well-being and mental health. In this context, the present study examined the effectiveness of a positive psychology intervention, following Seligman's guidelines (2002), on protective factors against psychological distress (emotional intelligence, resilience, interpersonal skills) in a sample of 45 adolescents aged between 13 and 16 years. All participants completed a battery of standardized questionnaires before (T0) and after (T1) the positive psychology intervention. Overall, the results show significant improvements in the scores on questionnaires assessing skills related to emotional intelligence and interpersonal competencies, highlighting how positive psychology can make adolescents more able to recognizing, understanding, and managing their own and others' emotions, with significant reductions in emotional distress experienced in social situations. Positive psychology interventions in adolescence contribute to strengthening protective psychological aspects, thus proving useful in preventing psychological distress.

Key words

Positive psychology; Psychological well-being; Adolescence; Quality of life

Introduzione

L'adolescenza è una fase critica dello sviluppo, caratterizzata da significativi cambiamenti fisici, cognitivi e socio-emotivi (Steinberg, 2005). Queste trasformazioni possono avere importanti conseguenze sulla salute mentale e sul benessere psicologico (Kessler et al., 2001; Steinberg, 2005; Patel et al., 2007). Secondo un rapporto del Fondo delle Nazioni Unite per l'infanzia (UNICEF, 2021), un adolescente su sette tra i 10 e i 19 anni soffre di un disturbo mentale, e tra questi i più comuni sono i disturbi d'ansia e dell'umore (circa il 40%). In questo contesto, l'implementazione di interventi che promuovano il benessere psicologico, attraverso lo sviluppo adeguato di abilità emotive e sociali, risulta cruciale per avere un'adeguata salute mentale e prevenire il disagio psichico. In quest'ottica, negli ultimi anni, è emerso un approccio conosciuto come "Psicologia Positiva" (Seligman, 2002) che, non limitandosi ai sintomi della sofferenza psicologica e della psicopatologia, mira a costruire e potenziare i punti di forza e le risorse di un individuo con uno sguardo positivo rivolto verso la rappresentazione di sé, del mondo che ci circonda, e del futuro.

Ad oggi, sono disponibili diverse tipologie di interventi nell'ambito della Psicologia Positiva, riconducibili a due approcci principali: unidimensionali o multidimensionali (Owens e Waters, 2020; Tejada-Gallardo et al., 2020). In particolare, gli interventi di Psicologia Positiva unidimensionali, si concentrano su singoli esercizi volti a promuovere specifiche dimensioni del benessere, come la gratitudine, la speranza o i punti di forza del carattere (Froh et al., 2009; Marques et al., 2011; Proctor et al., 2011), mentre gli interventi multidimensionali si basano su esercizi che mirano simultaneamente a promuovere più dimensioni del benessere. Diversi studi (Rusk et al., 2017; Hendriks et al., 2020) suggeriscono che gli interventi multidimensionali, attraverso l'utilizzo di un programma integrato, possano offrire benefici duraturi sul benessere psicologico e ridurre il rischio di ricadute rispetto a quelli monodimensionali. Nell'ambito degli approcci multidimensionali di Psicologia Positiva l'intervento proposto da Seligman e colleghi (2009), emerge come una alternativa promettente, in quanto non solo si concentra sulla riduzione del disagio e sulla promozione del benessere psicologico ma risulta, inoltre definito sulla base di un framework metodologico chiaro e standardizzato.

Di fatti, tale approccio si incentra sulla teoria del benessere, conosciuta come 'PERMA' (Seligman e Csikszentmihalyi, 2000) che identifica cinque elementi fondamentali per incrementare il benessere psicologico, che sono: i. "Positive Emotions" ovvero le emozioni positive e l'esperienza di sentimenti piacevoli; ii. "Engagement", ovvero l'impegno che si prova quando si è completamente immersi in un'attività; iii. "Relationships", le relazioni positive e la presenza di legami sociali forti e significativi; iv. "Meaning" inteso come significato o meglio la sensazione che la propria vita abbia uno scopo e che le proprie azioni siano significative; v. "Accomplishment" e in particolare il senso di riuscita e di raggiungimento di obiettivi personali e professionali che portano a un senso di successo e di competenza (Seligman e Csikszentmihalyi, 2000). La teoria di Seligman sottolinea l'importanza di bilanciare questi cinque elementi per raggiungere un alto livello di benessere, di resilienza e realizzazione personale (Seligman et al., 2005). Il corretto equilibrio di queste abilità conduce ad una maggiore soddisfazione della propria vita, e avrebbe un importante ruolo nella prevenzione dei disturbi mentali (Wood et al., 2010). In questo contesto un elemento chiave è l'autostima, intesa come la valutazione positiva di sé e delle proprie capacità, e si rivela fondamentale nel processo di costruzione del benessere psicologico. Infatti, è stato mostrato che individui che hanno sviluppato un buon livello di benessere psicologico attraverso alti livelli di autostima e di fiducia in sé stessi e nelle proprie capacità cognitive, emotive, ed interpersonali, sono più capaci di regolare le proprie emozioni quando devono affrontare situazioni stressanti (Blanchard-Fields e Coats, 2008).

In particolare, gli interventi di Psicologia Positiva multidimensionali che mirano a consolidare costrutti come la gratitudine, la resilienza, la speranza, le emozioni positive, e i propri punti di forza, hanno un effetto specifico sui livelli di emozioni positive sperimentate quotidianamente e sulla riduzione della frequenza dei pensieri negativi (Seligman et al., 2005; Carson et al., 2012), migliorando il benessere psicologico complessivo degli individui (Seligman et al., 2005; Carson et al., 2012).

Così, la riduzione di pensieri negativi accompagnata dall'aumento di emozioni positive nella quotidianità può essere utile anche in età adolescenziale agendo positivamente sul rendimento scolastico e sulle relazioni con i pari (Seligman et al., 2009; Green et al., 2011).

Pertanto, l'obiettivo del presente studio è valutare l'efficacia di un intervento multidimensionale di Psicologia Positiva su un campione di adolescenti, ipotizzando che tale intervento possa potenziare le diverse dimensioni del benessere psicologico, tra cui l'intelligenza emotiva, la resilienza e le competenze interpersonali.

Metodi

Partecipanti

In questo studio sono stati reclutati settantadue adolescenti di età compresa tra i 13 e i 16 anni (32 maschi e 13 femmine) con sviluppo tipico, presso l'Istituto Superiore Istruzione Secondaria Francesco De Sanctis – Oscar D'Agostino di Avellino (Italia). Sono state coinvolte sei classi dei primi tre anni di studio, che hanno partecipato singolarmente ad ogni incontro dell'intervento di Psicologia Positiva. Tutti i partecipanti avevano nazionalità italiana, ed assenza di disturbi psichiatrici e cognitivi, hanno preso parte

allo studio in modo volontario, e per tutti è stato ottenuto il consenso informato scritto dai genitori o dai tutori legali.

Lo studio è stato approvato dal comitato etico locale ed è stato condotto secondo gli standard etici stabiliti dalla Dichiarazione di Helsinki del 1964.

Procedura

Il progetto è stato realizzato in collaborazione con il “Rotary Club Avellino” - Distretto 2101, che ha svolto il ruolo di finanziatore e promotore dell'iniziativa, con l'intento di implementare strategie e programmi di prevenzione finalizzati a migliorare il benessere psicologico dei giovani.

Il presente studio è stato articolato in due fasi distinte. Nella prima fase (marzo/aprile 2024), è stata condotta una valutazione iniziale (T0), durante la quale tutti i partecipanti hanno completato una batteria di questionari standardizzati, selezionati per misurare lo stato di salute mentale, il benessere emotivo, e il profilo comportamentale. I questionari sono stati compilati attraverso la piattaforma Google Moduli. Per facilitare l'accesso ad essi è stato fornito un QR code che indirizzava direttamente alla pagina di compilazione. Questo metodo ha garantito l'anonimato dei partecipanti, in quanto nessuna informazione personale identificabile è stata raccolta durante la compilazione.

La valutazione è stata effettuata in classe, progettato per garantire tranquillità e comfort, affinché i partecipanti potessero rispondere in modo sincero e privo di influenze esterne. Dopo una settimana dalla valutazione iniziale, tutti i partecipanti hanno preso parte a dieci incontri di psicoeducazione incentrati sul modello “PERMA” di Seligman e Csikszentmihalyi (2000), condotti da psicologi con una formazione specifica sulla Psicologia Positiva. Gli incontri si sono svolti con frequenza bi-settimanale, consentendo un approfondimento graduale dei temi trattati, sempre all'interno delle singole classi.

Alla conclusione di ciascun incontro, è stato riservato un momento per l'ascolto attivo, in cui i partecipanti hanno avuto l'opportunità di esprimere le proprie opinioni e riflessioni sugli argomenti discussi. Al termine degli interventi, due mesi dopo la valutazione al T0, tutti i partecipanti sono stati sottoposti nuovamente alla stessa batteria di questionari (T1) già somministrata in precedenza.

Infine, è stato organizzato un incontro che ha coinvolto tutte le classi e tutti gli psicologi partecipanti al progetto. Durante questo incontro, sono stati presentati i risultati ottenuti, con particolare attenzione sull'importanza di promuovere il benessere psicologico durante l'adolescenza. Questa condivisione ha offerto l'opportunità di discutere le implicazioni dei risultati e di sottolineare il ruolo cruciale del supporto psicologico in questa fase di vita.

Strumenti

I singoli strumenti sono stati selezionati per garantire una valutazione il più possibile coerente e in relazione al modello “PERMA” di Seligman e Csikszentmihalyi (2000), al fine di quantificare l'effetto dell'intervento psicoeducativo e poter valutare eventuali miglioramenti nel benessere psicologico nei partecipanti.

Nel dettaglio la batteria di questionari standardizzati somministrati comprendeva:

- Il Questionario sulla salute del paziente per adolescenti 9-item (PHQ-9; Johnson et al., 2002), uno strumento di auto-somministrazione che valuta la presenza e la gravità dei sintomi

depressivi nei giovani. Il questionario comprende 9 items basati sui criteri del DSM-5 (American Psychiatric Association, 2003) per il disturbo depressivo maggiore, che esplorano la frequenza dei sintomi depressivi nelle ultime due settimane. Esso ha una buona consistenza interna (alpha di Cronbach: 0,86-0,89; Kroenke et al., 2001) e le risposte vengono date su una scala Likert a quattro punti che varia da 0 (per niente) a 3 (quasi ogni giorno). Il punteggio totale varia da 0 a 27, con punteggi più alti indicativi di una maggiore gravità dei sintomi depressivi;

- La Scala dell'Intelligenza Emotiva (EIS; Schutte et al., 1998; Grazzani et al., 2009), uno strumento di auto-somministrazione progettato per misurare l'intelligenza emotiva, definita come la capacità di riconoscere, comprendere, gestire e utilizzare in modo efficace le proprie emozioni e quelle altrui. La scala è composta da 33 item, ogni item è valutato su una scala Likert a cinque punti che varia da 1 (per niente d'accordo) a 5 (totalmente d'accordo). Il punteggio totale va da un range di 33 a 165, con punteggi più alti indicativi di un maggiore livello di competenza in ciascuna dimensione dell'intelligenza emotiva. La scala include le seguenti sottoscale: Valutazione ed Espressione delle Emozioni Riferite agli Altri (alpha di Cronbach:0,73; Grazzani et al., 2009), che misura la capacità di un individuo di percepire e comprendere accuratamente le emozioni degli altri; Valutazione ed Espressione delle Emozioni Riferite a Sé (alpha di Cronbach:0,66; Grazzani et al., 2009), valuta la capacità di un individuo di riconoscere e comprendere le proprie emozioni; Regolazione ed Uso delle Emozioni/Umore (alpha di Cronbach:0,72; Grazzani et al., 2009), misura la capacità di gestire e regolare le proprie emozioni in modo efficace.

- La Scala per la resilienza 14-Item (RS-14; Wagnild, 2009; Callegari et al., 2016), uno strumento di auto-somministrazione progettato per misurare la resilienza, definita come la capacità di affrontare lo stress e le difficoltà in modo adattivo. La scala ha una buona consistenza interna (alpha di Cronbach: 0,88; Callegari et al., 2016), ed è composta da 14 item, ogni item è valutato su una scala Likert a sette punti, che vanno da 1 (completamente in disaccordo) a 7 (completamente d'accordo). Il punteggio totale della RS-14 va da un range di 14 a 98, con punteggi più alti indicativi di un maggiore livello di resilienza.

- La Scala per il comportamento interpersonale (SIB; Arrindell et al., 2002), una scala di auto-somministrazione che valuta i comportamenti interpersonali e assertivi. Essa è composta da 25 item in cui viene richiesto di indicare il livello di disagio provato nel mettere in atto quel comportamento (alfa di Cronbach: 0,90; Arrindell et al., 2002) e la probabilità di comportarsi secondo il modo descritto comportamento (alfa di Cronbach: 0,85; Arrindell et al., 2002). La scala consente di esplorare diverse dimensioni del comportamento interpersonale, quali: Assertività Positiva, ovvero la capacità di dare e ricevere lodi o complimenti; Assertività Negativa ovvero, la capacità di rifiutare le richieste in difesa dei propri diritti in una situazione pubblica; Assertività Sociale, ovvero la capacità di esprimere le proprie opinioni e iniziare una conversazione con estranei; Espressione e Gestione dei Limiti Personali ovvero la capacità di riconoscere il proprio fallimento o le proprie carenze, saper gestire le critiche e le pressioni e chiedere aiuto. Il punteggio totale può variare da un minimo di 0 a un massimo di 125, punteggi più alti riflettono una maggiore competenza sociale e una migliore capacità di interazione.

- L'Organizzazione Mondiale della Sanità sullo Stato di Benessere (OMS-5; World Health Organization, 1998), una misura auto-valutativa progettata per valutare il benessere psicologico generale degli individui. Il questionario ha una buona consistenza interna (alfa di Cronbach: 0,79-0,91; Nicolucci et al., 2004), ed è composto da 5 item, ognuno dei quali misura un aspetto specifico del benessere positivo. Ogni item viene valutato su una scala Likert a 6 punti, che varia da 0 (per niente) a 5 (sempre). La somma dei punteggi di questi item fornisce un'indicazione complessiva del livello di benessere psicologico, con punteggi più alti indicanti un maggiore benessere.

Intervento

L'intervento di Psicologia Positiva è stato condotto attraverso 10 incontri di psicoeducazione seguendo il modello "PERMA" di Seligman e Csikszentmihalyi (2000). Il protocollo clinico prevedeva un primo incontro sull'Introduzione alla Psicologia Positiva; un secondo incontro volto a determinare i punti di forza del carattere e a consolidarli; un terzo incontro per favorire la felicità e il benessere nella vita quotidiana, un quarto incontro incentrato sul senso della vita e sull'autorealizzazione; un quinto incontro per descrivere e conoscere le 10 emozioni positive; un sesto incontro per definire e lavorare sull'autostima e la fiducia in sé stessi; un settimo incontro per definire il concetto di ottimismo e per lavorare sulla motivazione; un ottavo incontro per imparare a gestire più facilmente i propri pensieri al fine di poter essere presente nel momento smettendo; un nono incontro per rafforzare, sviluppare e mantenere nel tempo le relazioni sociali; ed infine, un decimo incontro di fine programma volto a prendere coscienza dei progressi fatti. Per maggiori dettagli sugli obiettivi e le attività svolte durante gli incontri vedi Tabella 1.

Tabella 1. Descrizione delle attività svolte durante i 10 incontri di psicoeducazione basati sull'approccio della Psicologia Positiva

Numero Incontro	Titolo Incontro	Obiettivo	Descrizione dell'attività proposta
1	Introduzione alla Psicologia Positiva	L'obiettivo è quello di mostrare come la Psicologia Positiva mira a migliorare il benessere quotidiano basandosi sui 5 pilastri: le emozioni positive, l'impegno, le relazioni umane, il senso e lo scopo della vita, la realizzazione (Seligman, 2000). Nello specifico provare emozioni positive, godere dei piccoli momenti è importante per condurre una vita piacevole. L'impegno permettere di dedicarsi completamente ad un'attività utilizzando i propri punti di forza. Il senso e lo scopo della vita permettono di condurre una vita piena di significato che si basa anche sulle relazioni interpersonali.	Attraverso l'ausilio di una presentazione PowerPoint, sono stati presentati in modo interattivo i concetti fondamentali della Psicologia Positiva. Successivamente, è stato dedicato un momento a un'attività interattiva, durante la quale i partecipanti sono stati invitati a descrivere e discutere le seguenti domande: <ul style="list-style-type: none"> - Pensa al momento, all'azione o alla situazione che ti ha dato più soddisfazione e benessere negli ultimi giorni. - In questo preciso momento, cosa provi? Felicità? Stress? Stanchezza? Ansia? Descrivi con parole o aggettivi semplici ciò che senti nel qui e ora. - Quali azioni, attività o eventi hai in programma per i prossimi giorni che già solo a pensarci ti procurano felicità ed entusiasmo? - Cita un'attività che ti piace fare e che ti dà un'emozione positiva autentica. - Descrivi un'attività o un'occupazione che ti procuri una sensazione di "flow". - Cita un'attività in cui ti senti coinvolto e che ti fa sentire utile.

2	Determinare i punti di forza del carattere e consolidarli	L'obiettivo è quello di definire e potenziare i 24 punti di forza del carattere che definiscono le proprie sei virtù (saggezza, coraggio, umanità, giustizia, moderazione/temperanza e trascendenza/spiritualità) affinché vengano sfruttati al meglio nel quotidiano e nelle situazioni della vita reale.	<p>Nella fase iniziale dell'incontro è stata presentata una dettagliata esposizione PowerPoint che ha illustrato i 24 punti di forza, evidenziando come questi si suddividano nelle sei virtù fondamentali. Successivamente, i partecipanti hanno svolto esercizi pratici volti a riflettere sui propri punti di forza. Questi esercizi hanno incoraggiato una concentrazione sui tratti caratteriali individuali, aiutando a sviluppare una maggiore consapevolezza delle proprie capacità e come queste possano essere utilizzate per promuovere il benessere personale.</p> <p>I partecipanti hanno svolto la seguente attività pratica:</p> <ul style="list-style-type: none"> - Identificazione dei Punti di Forza: Prenditi del tempo per riflettere sui tuoi punti di forza personali. Pensa a situazioni in cui hai avuto successo o in cui hai ricevuto complimenti. Se hai difficoltà a identificarli, puoi chiedere a persone di fiducia, come amici, familiari o colleghi, di aiutarti. - Elenco dei Punti di Forza: Crea una lista dei tuoi punti di forza. Cerca di essere specifico e dettagliato. - Riflessione Quotidiana: Ogni giorno, scegli uno dei tuoi punti di forza dalla lista. Scrivi un breve paragrafo su come hai utilizzato questo punto di forza in una situazione recente. Descrivi la situazione, come hai messo in pratica il tuo punto di forza, e quale impatto ha avuto sul risultato. - Riflessione Finale: Dopo aver riflettuto su un punto di forza, prenditi un momento per riconoscere come questo ti ha aiutato e come ti fa sentire. Nota se questa pratica influisce sul tuo umore e sulla tua autostima nel corso del tempo.
3	Felicità e benessere: trovare e avere positività nella vita quotidiana	L'obiettivo è comprendere la nascita della psicologia positiva facendo una distinzione tra felicità (costrutto reale, quantificabile) e benessere (costrutto astratto, non quantificabile), e comprendere come le emozioni positive costituiscono uno dei legami con la felicità più evidenti e tangibili.	<p>Nella prima parte dell'incontro, è stata presentata una spiegazione dettagliata, supportata da una presentazione PowerPoint, sull'importanza delle emozioni positive e sul loro ruolo nel promuovere una visione ottimistica della vita quotidiana. Successivamente, i partecipanti hanno partecipato a un'attività pratica che li invitava a scrivere, alla fine di ogni giornata, tre eventi positivi che si erano verificati, siano essi grandi o piccoli. È stata enfatizzata l'importanza di specificare perché ciascuno di questi eventi fosse significativo, sottolineando che anche i piccoli gesti hanno un valore fondamentale. Questa pratica mira a sviluppare l'abitudine di apprezzare le piccole cose della vita quotidiana. La riflessione finale su un aspetto positivo della giornata, anche se di modesta entità, può infatti trasformare la propria prospettiva e arricchire l'esperienza di vita.</p>
4	Senso della vita e autorealizzazione, definizione degli obiettivi e relativo raggiungimento	L'obiettivo è definire le nozioni del senso della vita e dell'appagamento, e stabilire una gerarchia di obiettivi attraverso il metodo SMART (ogni obiettivo deve essere specifico, misurabile, accessibile, realistico, temporizzabile), che potrebbe contribuire alla ricerca della felicità e del benessere.	<p>Nella prima parte dell'incontro, è stata fornita un'analisi approfondita, supportata da una presentazione PowerPoint, sui concetti di relazione personale e senso della vita, evidenziando come questi elementi possano influenzare l'autorealizzazione. In questo contesto, è stato sottolineato che fissare obiettivi personali può contribuire a creare un senso di controllo sulla propria esistenza, a incrementare l'autostima e a conferire significato alla vita.</p> <p>Nella seconda parte, caratterizzata da un approccio più interattivo, ai partecipanti è stato proposto l'esercizio "La Lista dei Desideri". Questo esercizio prevedeva che scrivessero i propri desideri, li classificassero gerarchicamente in base a quelli di maggiore priorità e, successivamente, valutassero la loro realizzabilità. Questa attività ha permesso ai partecipanti di riflettere attivamente sui propri obiettivi e di sviluppare una maggiore consapevolezza delle proprie aspirazioni.</p>

5	Le emozioni positive	L'obiettivo è descrivere e conoscere le 10 emozioni positive (gioia, gratitudine, serenità, curiosità, speranza, fierezza, divertimento, ispirazione, ammirazione, amore), e conoscere quali potrebbero essere le attività che suscitano emozioni positive (aiutare, comunicare con gli altri, giocare e divertirsi, imparare e fare attività fisica).	Nella prima parte dell'incontro, è stata presentata, tramite una presentazione PowerPoint, un'analisi delle 10 emozioni positive e del loro significativo impatto sul benessere e sulla felicità quotidiana. È stata enfatizzata l'importanza di coltivare comportamenti che facilitino l'esperienza di emozioni positive, evidenziando come il pensiero positivo possa influenzare le azioni quotidiane. È stata anche illustrata la differenza tra emozioni positive e pensieri negativi, questi ultimi spesso associati a paure e ansie. Nella seconda parte dell'incontro, sono state svolte diverse attività interattive. Nella prima, ai partecipanti è stato chiesto di identificare le 10 emozioni positive che sperimentavano meno frequentemente e quelle che avrebbero voluto provare più spesso, scegliendo da un elenco di categorie presentate. Successivamente, è stato richiesto di riflettere su queste emozioni e su come potessero aumentare le opportunità di viverle nella vita quotidiana. Un'ulteriore attività si è concentrata sulla consapevolezza dell'impatto dei propri giudizi e pensieri negativi sulle emozioni personali. I partecipanti sono stati invitati a identificare e descrivere i giudizi negativi che avevano emesso durante la settimana, sia nei confronti degli altri che di se stessi, in diverse situazioni. Hanno quindi analizzato ciascun giudizio, la situazione che lo aveva provocato e le emozioni associate, con l'obiettivo di trasformare il giudizio negativo in un pensiero più realistico e positivo.
6	Lavoro sull'autostima e sulla fiducia in sé stessi	L'obiettivo è definire concetti come l'autostima e la fiducia in sé stessi, capire la differenza e il legame tra questi due concetti e come questi hanno un impatto sulla qualità di vita.	Nella prima parte dell'incontro, è stata presentata, attraverso una presentazione PowerPoint, una panoramica approfondita sull'autostima e sul suo significativo impatto sulla qualità della vita. È stato evidenziato come l'autostima influisca positivamente sulla fiducia in sé stessi, mostrando il forte legame tra questi due costrutti psicologici. La relazione tra autostima e fiducia è stata illustrata attraverso esempi pratici, che hanno reso chiaro come un miglioramento in una di queste aree possa portare a progressi nell'altra. Nella seconda parte dell'incontro, sono state proposte in modo interattivo diverse strategie per potenziare sia l'autostima che la fiducia in sé. Tra queste, è stata sottolineata l'importanza di prendere consapevolezza delle situazioni e dei pensieri negativi che influenzano la propria vita quotidiana. I partecipanti sono stati invitati a riflettere sui momenti in cui si giudicano o si criticano, ponendosi domande chiave: quali sono le origini di queste valutazioni? Su quali basi si fondano i loro commenti auto-critici? Perché tendono a minimizzare le proprie capacità? Questa riflessione ha lo scopo di aiutare i partecipanti a identificare e contestare i pensieri negativi, promuovendo una visione più realistica e positiva di sé stessi.
7	Ottimismo e motivazione	L'obiettivo è definire i concetti di ottimismo e di motivazione e conoscere gli strumenti che permetteranno di coltivare l'ottimismo e la motivazione.	Nella prima parte dell'incontro, è stata presentata, attraverso una presentazione PowerPoint, un'analisi approfondita dell'ottimismo e dei molteplici vantaggi che esso offre nel raggiungimento degli obiettivi personali. È stata chiarita la distinzione tra ottimismo e pessimismo, evidenziando come quest'ultimo possa avere effetti permanenti su diversi aspetti della vita quotidiana. Successivamente, è stato affrontato il concetto di motivazione, esplorando come essa spinga l'individuo ad intraprendere azioni e a perseverare fino al conseguimento dei propri traguardi. Nella seconda parte dell'incontro, sono state proposte attività più dinamiche, caratterizzate da discussioni attive, con l'obiettivo di identificare strategie per dissipare il pessimismo e coltivare l'ottimismo. Le strategie discusse includevano: <ul style="list-style-type: none"> - Smettere di rimuginare sul passato - Riconoscere la propria responsabilità negli eventi positivi - Sfruttare i punti di forza del proprio carattere - Imparare a vedere il mondo in modo positivo - Ridurre i pensieri negativi

			<ul style="list-style-type: none"> - Circondarsi di persone positive e ottimiste - Superare le convinzioni limitanti e la paura del fallimento <p>Inoltre, gli esercizi mirati a mantenere alta la motivazione si basavano sui seguenti aspetti:</p> <ul style="list-style-type: none"> - Prendere consapevolezza delle attività che costituiscono fonte di motivazione - Analizzare i costi e i benefici di comportamenti inadeguati - Non rimandare a domani ciò che si può fare oggi - Utilizzare liste di attività da completare (to-do list) - Evitare di parlare dei propri obiettivi al condizionale - Prendersi cura di sé in generale - Riadattare regolarmente i propri obiettivi <p>Queste attività hanno permesso di incoraggiare una riflessione profonda e di promuovere un approccio proattivo verso il miglioramento del benessere psicologico.</p> <p>Nella prima parte dell'incontro, è stata presentata, mediante una presentazione PowerPoint, un'analisi approfondita sull'importanza di esaminare i propri pensieri in modo oggettivo. Questo processo mira a valutare se i pensieri che affollano la mente siano gradevoli o sgradevoli.</p> <p>Nella fase successiva, è stata introdotta un'attività pratica che ha invitato i partecipanti a riflettere sull'impatto dei propri pensieri sulle emozioni. È stato sottolineato come i pensieri abbiano il potere di plasmare i sentimenti, influenzandoli sia in modo positivo che negativo.</p> <p>Diventando consapevoli del legame tra pensieri ed emozioni, i partecipanti possono apprendere strategie per interrompere cicli negativi o avviare spirali di pensieri e sentimenti positivi. Questa consapevolezza rappresenta un passo fondamentale verso il miglioramento del benessere psicologico.</p> <p>Nella prima parte dell'incontro, è stata presentata, tramite una presentazione PowerPoint, un'analisi approfondita su uno dei pilastri fondamentali della psicologia positiva: le relazioni sociali, che rivestono un ruolo cruciale nella ricerca della felicità e del benessere. È stato evidenziato come tali relazioni possano essere rafforzate attraverso costrutti come la gratitudine e la comunicazione. La gratitudine, infatti, è un'emozione positiva che favorisce lo sviluppo, la creazione e il miglioramento delle relazioni sociali, mentre una comunicazione efficace è essenziale per instaurare relazioni positive.</p> <p>Nella seconda parte dell'incontro, sono stati condotti esercizi pratici, tra cui la "lettera della gratitudine", che invita i partecipanti a scrivere a una persona che ha avuto un impatto significativo sulla propria vita, esprimendo la riconoscenza per il suo supporto. Un altro esercizio proposto è stato il "diario della gratitudine", in cui ai partecipanti è stato chiesto di annotare quotidianamente le esperienze positive, incluse azioni che hanno generato un senso di realizzazione, momenti di piacere derivanti dall'aiuto offerto ad altri, e semplici gioie sensoriali come ciò che hanno visto, sentito, assaporato, toccato o annusato.</p> <p>In questo incontro finale, oltre a condurre una valutazione per esaminare gli effetti dell'intervento, sono stati forniti strumenti pratici per monitorare i progressi ottenuti. Sono stati anche condivisi consigli pratici riguardanti esercizi da svolgere quotidianamente, che possono avere un impatto positivo sul benessere psicologico.</p> <p>Un esempio è la pratica della cura di sé, che implica dedicare tempo e attenzione al proprio benessere fisico, emotivo e mentale. Questo comprende l'auto-compassione, ovvero il praticare un dialogo interno caratterizzato da gentilezza e comprensione, specialmente nei momenti di difficoltà.</p> <p>Un altro strumento efficace è la riflessione sui successi</p>
8	Istante presente e piena coscienza	L'obiettivo è imparare a conoscere, ascoltare, fermare, analizzare i propri pensieri, per poi includerli se necessario, nelle proprie azioni.	
9	Comunicazione, gratitudine e relazioni sociali	L'obiettivo è comprendere come rafforzarle, svilupparle e mantenerle nel tempo le relazioni sociali, e come queste possono essere rafforzate da costrutti come la gratitudine e la comunicazione.	
10	Valutazione di fine programma e consigli per le pratiche di coaching	L'obiettivo è quello di effettuare una valutazione di fine programma, e aiutare i partecipanti a prendere coscienza dei progressi fatti.	

quotidiani e settimanali, che aiuta a riconoscere e apprezzare i propri progressi e contributi. Infine, la creazione di una routine positiva per iniziare e concludere la giornata lavorativa può avere un impatto significativo sulla produttività e sul benessere mentale.

Analisi Statistica

La normalità della distribuzione è stata valutata utilizzando il test di Kolmogorov-Smirnov, condotto sui punteggi ottenuti ai questionari somministrati prima dell'intervento di Psicologia Positiva. Dai risultati, è emerso che la maggior parte delle variabili risultavano normalmente distribuite (PHQ: $D=0,166$; $p=0,003$; EIS: $D=0,097$; $p=0,200$; RS-14: $D=0,113$, $p=0,199$; SIB totale disagio: $D=0,090$; $p=0,200$; SIB totale probabilità: $D=0,091$; $p=0,189$; OMS: $D=0,112$; $p=0,199$) e per tale ragione sono stati eseguite analisi parametriche.

È stata utilizzata l'analisi multivariata (MANOVA) per misure ripetute, con una variabile indipendente a due livelli: il tempo, considerando i punteggi prima (T0) e dopo (T1) l'intervento; e come variabili dipendenti i punteggi ottenuti alla batteria di questionari somministrati.

Le analisi statistiche sono state condotte utilizzando SPSS-21 (SPSS lic, Chiago. I11), con un valore $p < 0,05$ considerato statisticamente significativo.

Risultati

Un totale di 72 partecipanti ha compilato la batteria di valutazione prima dell'intervento di Psicologia Positiva (T0). Tuttavia, solo 45 partecipanti (32 maschi e 13 femmine, età media: $14,80 \pm 0,67$) erano presenti durante la valutazione dopo l'intervento (T1).

L'analisi della MANOVA a misure ripetute ha mostrato un effetto statisticamente significativo del tempo ($F=2,805$; $p=0,009$). Nello specifico, è emerso un effetto significativo sui punteggi ottenuti al questionario che valuta l'intelligenza emotiva (EIS; $F=22,171$ $p \leq 0,001$), in particolare nelle sottoscale Valutazione ed Espressione delle emozioni riferite agli altri ($F=16,640$; $p \leq 0,001$), e Emozioni riferite a sé ($F=19,033$; $p \leq 0,001$); e nel questionario che valuta i comportamenti interpersonali ($F=5,254$, $p=0,027$), in particolare, nelle sottoscale Assertività Positiva ($F=4,336$; $p=0,042$); Assertività Negativa ($F=6,544$; $p=0,014$); e Assertività Sociale ($F=7,828$; $p=0,008$; Tabella 2).

Non emergono differenze statisticamente significative nei punteggi ottenuti agli altri questionari somministrati in seguito all'intervento ($F=2,444$, $p>0,125$). Medie e deviazioni standard dei punteggi ottenuti ai questionari prima e dopo l'intervento di Psicologia Positiva sono riportati in Tabella 2.

Tabella 2. Medie e deviazioni standard dei punteggi ottenuti ai questionari somministrati prima (T0) e dopo (T1) l'intervento di Psicologia Positiva

Variabili Comportamentali		Partecipanti (n=45)			
		M±DS	Min-Max	F	p
PHQ-9	T0	6,64±3,89	0-16	2,444	0,125
	T1	7,31±4,34	0-16		
EIS Totale	T0	108,84±14,12	76-131	22,171	≤0,001
	T1	115,53±11,91	89-135		
EIS- valutazione ed espressione delle emozioni riferite agli altri	T0	32,49±4,70	22-40	16,640	≤0,001
	T1	35,04±4,64	23-44		
EIS- valutazione ed espressione delle emozioni riferite a sé	T0	13±3,25	6-17	19,033	≤0,001
	T1	14,49±3,20	6-20		
EIS- Regolazione ed uso delle emozioni	T0	32,22±4,18	24-40	2,216	0,144
	T1	31,27±4,71	22-40		
SIB-disagio Totale	T0	63,44±17,81	27-105	5,254	0,027
	T1	57,24±15,85	25-100		
SIB-disagio assertività positiva	T0	16,60±5,63	6-30	4,336	0,042
	T1	15,02±5,42	6-30		
SIB-disagio assertività negativa	T0	17,64±5,78	7-33	6,544	0,014
	T1	15,49±5,06	7-34		
SIB- disagio espressione e gestione dei limiti personali	T0	13,42±5,07	6-30	0,127	0,723
	T1	13,09±4,60	6-30		
SIB- disagio assertività sociale	T0	15,78±5,10	6-30	7,828	0,008
	T1	13,64±3,89	6-30		
SIB-SF-probabilità Totale	T0	70,84±12,31	36-89	0,039	0,844
	T1	70,44±16,70	25-112		
RS-14 Totale	T0	61,47±15,22	14-89	1,314	0,258
	T1	59,11±15,74	14-95		
OMS-5	T0	11,98±5,44	1-25	2,191	0,146
	T1	13,33±5,15	2-25		

Note: T0, pre-intervento; T1, post-intervento; EIS, Scala per l'Intelligenza Emotiva; SIB, Scala per il Comportamento Interpersonale; PHQ-9, Questionario sulla salute del paziente per adolescenti 9-item; RS-14, Scala per la resilienza 14-Item; OMS-5, Questionario dell'Organizzazione Mondiale della Sanità sullo stato di benessere.

In grassetto sono riportate le differenze statisticamente significative p<0,05

Discussione

L'obiettivo del presente studio era quello di valutare l'efficacia di un intervento multidimensionale di Psicologia Positiva sul benessere psicologico degli adolescenti. Dai risultati è emerso che, a seguito dell'intervento, i partecipanti hanno mostrato un significativo miglioramento delle competenze relative all'intelligenza emotiva e alle abilità interpersonali. In particolare, è stata osservata una maggiore capacità di riconoscere, comprendere e comunicare le proprie emozioni e quelle degli altri, il che ha portato a una riduzione del disagio percepito in situazioni interpersonali complesse, in cui è richiesto di affermare i propri diritti o gestire emozioni difficili come rabbia e frustrazione.

Tali risultati indicano che un intervento multidimensionale basato sulla Psicologia Positiva potrebbe fornire agli adolescenti una serie di strumenti pratici utili per migliorare il loro benessere psicologico. Tecniche come il diario della gratitudine e la meditazione, già ben documentate nella letteratura come efficaci per aumentare il benessere emotivo (Emmons e McCullough, 2003; Fredrickson, 2001), possono essere integrate in un vero e proprio "prontuario" di strategie pratiche. Questo prontuario permetterebbe agli adolescenti di acquisire competenze per una gestione più efficace delle proprie emozioni, migliorando così anche le loro relazioni sociali.

Tuttavia, è importante sottolineare che, sebbene gli interventi di Psicologia Positiva abbiano un impatto positivo nel potenziamento delle emozioni positive, come la gratitudine e la speranza, la loro efficacia nel migliorare la regolazione delle emozioni negative potrebbe essere limitata. La regolazione emotiva, soprattutto quando si tratta di emozioni negative intense o persistenti, spesso richiede approcci terapeutici più mirati, come la Terapia Cognitivo Comportamentale (Halder e Mahato, 2019). Questo suggerisce che l'intervento di Psicologia Positiva potrebbe essere utilizzato in combinazione con altre forme di trattamento, per ottenere un miglioramento più completo delle competenze emotive.

Inoltre, va considerato che lo sviluppo della resilienza e del benessere psicologico negli adolescenti è fortemente influenzato dal contesto familiare e sociale in cui vivono. Alcuni studi hanno mostrato che il sostegno familiare e il clima sociale possono influire significativamente sull'efficacia degli interventi psicologici (Cao et al., 2024), evidenziando come l'intervento di Psicologia Positiva potrebbe trarre maggiore beneficio se integrato con strategie di supporto che tengano conto del contesto in cui gli adolescenti si trovano.

Un altro aspetto da considerare è l'importanza della continuità dell'intervento. Come riportato da Lyubomirsky e colleghi. (2005), gli effetti positivi delle pratiche di Psicologia Positiva, come la gratitudine e la meditazione, tendono a intensificarsi con un uso regolare e prolungato nel tempo. Ciò implica che programmi di intervento a lungo termine potrebbero essere più efficaci nel promuovere un cambiamento duraturo nelle capacità emotive e nel benessere psicologico degli adolescenti.

Il presente studio non è privo di limitazioni. In primo luogo, la dimensione campionaria ridotta non permette di generalizzare pienamente i risultati a una popolazione più ampia di adolescenti. In secondo luogo, l'assenza di un gruppo di controllo limita la possibilità di stabilire un confronto diretto con altri approcci terapeutici o con l'assenza di intervento. Ulteriori studi dovrebbero mirare a replicare questi risultati su campioni più ampi e includere condizioni di controllo per poter meglio valutare l'efficacia comparativa della Psicologia Positiva rispetto ad altre forme di trattamento.

In sintesi, i risultati di questo studio preliminare suggeriscono che gli interventi di Psicologia Positiva possano essere uno strumento utile per migliorare il benessere psicologico e le competenze interpersonali degli adolescenti. Tuttavia, il loro utilizzo potrebbe essere ottimizzato combinandoli con approcci terapeutici più specifici per la regolazione emotiva, così come considerando l'influenza del contesto familiare e sociale per favorire un approccio più integrato e completo.

Il presente studio è stato concepito come un'analisi esplorativa iniziale per valutare l'efficacia dell'intervento di Psicologia Positiva negli adolescenti, pertanto, futuri studi dovrebbero prevedere l'uso di disegni sperimentali che includano un campione più ampio, gruppi di controllo e follow-up a lungo termine per valutare la persistenza degli effetti osservati. Ulteriori ricerche sono necessarie per esplorare l'utilizzo di interventi di psicoeducazione basati sulla Psicologia Positiva come strumento preventivo nel contesto adolescenziale.

Conclusioni

Gli interventi basati sulla Psicologia Positiva rappresentano strumenti pratici ed efficaci in quanto forniscono strategie utili e utilizzabili nella vita quotidiana per l'ottimizzazione dell'espressione emotiva, la riduzione del disagio correlato all'affermazione dei propri diritti, l'articolazione delle proprie ragioni e

la partecipazione attiva in contesti sociali. Incrementando la consapevolezza emotiva, l'autostima e le competenze sociali, tali interventi contribuiscono significativamente al miglioramento del benessere individuale e alla promozione di relazioni interpersonali più sane e soddisfacenti. I risultati di questo studio sottolineano l'importanza di adottare un approccio positivo come prevenzione per il mantenimento del benessere psicologico negli adolescenti.

Finanziamenti

Il presente studio è stato sostenuto con un finanziamento del Rotary Club Avellino e da una sovvenzione del Rotary International-Distretto 2101 A.R. 2023-2024, ottenuto dal Club.

Ringraziamenti

Gli autori ringraziano l'Istituto Tecnico Agrario "Francesco De Sanctis – Oscar D'Agostino" di Avellino, il Dirigente Scolastico, Caterini Pietro, e le sue collaboratrici, Santucci Luisa e Glave Costantina per la collaborazione a questa ricerca.

Bibliografia

- American Psychiatric Association, DSM-5 Task Force. (2013). *Diagnostic and statistical manual of mental disorders: DSM-5™ (5th ed.)*. American Psychiatric Publishing
- Arrindell, W. A., Sanavio, E., e Sica, C. (2002). *Introducing a short form version of the scale for interpersonal behavior (s-SIB) for use in Italy*. *Psicoterapia Cognitiva e Comportamentale*.
- Blanchard-Fields, F., e Coats, A. H. (2008). The experience of anger and sadness in everyday problems impacts age differences in emotion regulation. *Developmental Psychology*, 44, 1547-1556.
- Callegari, C., Bertù, L., Lucano, M., Ielmini, M., Braggio, E., e Vender, S. (2016). Reliability and validity of the Italian version of the 14-item Resilience Scale. *Psychological Reports*, 118, 623-642.
- Cao, F., Li, J., Xin, W., e Cai, N. (2024). Impact of social support on the resilience of youth: mediating effects of coping styles. *Frontiers in public health*, 1331813.
- Carson, J., Muir, M., Clark, S., Wakely, E., e Chander, A. (2012). Piloting a gratitude intervention in a community mental health team. *Groupwork*, 20, 73-87.
- Emmons, R. A., e McCullough, M. E. (2003). Counting blessings versus burdens: an experimental investigation of gratitude and subjective well-being in daily life. *Journal of personality and social psychology*, 84, 377.
- Fredrickson, B. L. (2001). The role of positive emotions in positive psychology: The broaden-and-build theory of positive emotions. *American psychologist*, 56, 218.
- Froh, J. J., Kashdan, T. B., Ozimkowski, K. M., e Miller, N. (2009). Who benefits the most from a gratitude intervention in children and adolescents? Examining positive affect as a moderator. *The Journal of Positive Psychology*, 4, 408-422.
- Grazzani, I., Ornaghi, V., e Piralli, F. (2009). La valutazione delle competenze socio-emotive in età prescolare: Proprietà psicometriche della versione italiana della Schutte Self-Report Emotional Intelligence Scale. *Rivista di Psicologia Clinica*, 1, 46-63.
- Green, S., Oades, L., e Robinson, P. (2011). Positive education: Creating flourishing students, staff and schools. *InPsych*, 33, 37-43.
- Halder, S., e Mahato, A. K. (2019). Cognitive Behavior Therapy for Children and Adolescents: Challenges and Gaps in Practice. *Indian journal of psychological medicine*, 41(3), 279-283.
- Hendriks, T., Schotanus-Dijkstra, M., Hassankhan, A., De Jong, J., e Bohlmeijer, E. (2020). The efficacy of multi-component positive psychology interventions: A systematic review and meta-analysis of randomized controlled trials. *Journal of happiness studies*, 21, 357-390.
- Johnson, J. G., Harris, E. S., Spitzer, R. L., e Williams, J. B. W. (2002). The Patient Health Questionnaire for Adolescents: Validation of an instrument for the assessment of mental disorders among adolescent primary care patients. *Journal of Adolescent Health*, 30, 196-204.
- Kessler, R. C., Avenevoli, S., e Ries Merikangas, K. (2001). Mood disorders in children and adolescents: an epidemiologic perspective. *Biological psychiatry*, 49, 1002-1014.
- Kroenke, K., Spitzer, R. L., e Williams, J. B. (2001). The PHQ-9: validity of a brief depression severity measure. *Journal of general internal medicine*, 16(9), 606-613.

- Lyubomirsky, S., Sheldon, K. M., e Schkade, D. (2005). Pursuing happiness: The architecture of sustainable change. *Review of general psychology*, 9, 111-131.
- Marques, S. C., Lopez, S. J., e Pais-Ribeiro, J. L. (2011). Building hope for the future?: A program to foster strengths in middle school students. *Journal of Happiness Studies*, 12, 139-152.
- Nicolucci, A., Giorgino, R., Cucinotta, D., Zoppini, G., Muggeo, M., Squatrito, S., Corsi, A., Lostia, S., Pappalardo, L., Benaduce, E., Girelli, A., Galeone, F., Maldonato, A., Perriello, G., Pata, P., Marra, G., e Coronel, G. A. (2004). Validation of the Italian version of the WHO-Well-Being Questionnaire (WHO-WBQ) and the WHO-Diabetes Treatment Satisfaction Questionnaire (WHO-DTSQ). *Diabetes, nutrition & metabolism*, 17(4), 235-243.
- Owens, R. L., e Waters, L. (2020). What does positive psychology tell us about early intervention and prevention with children and adolescents? A review of positive psychological interventions with young people. *The Journal of Positive Psychology*, 15(5), 588-597.
- Patel, V., Flisher, A. J., Hetrick, S., e McGorry, P. (2007). Mental health of young people: a global public-health challenge. *Lancet* (London, England), 369, 1302-1313.
- Proctor, C., Tsukayama, E., Wood, A. M., Maltby, J., Eades, J. F., e Linley, P. A. (2011). Strengths Gym: The impact of a character strengths-based intervention on the life satisfaction and wellbeing of adolescents. *The Journal of Positive Psychology*, 6, 377-388.
- Rusk, R. D., Vella-Brodrick, D. A., e Waters, L. (2017). A complex dynamic systems approach to lasting positive change: The Synergistic Change Model. *The Journal of Positive Psychology*, 13, 406-418.
- Schutte, N. S., Malouff, J. M., Hall, L. E., Haggerty, D. J., Cooper, J. T., Golden, C. J., e Dornheim, L. (1998). Development and validation of a measure of emotional intelligence. *Personality and Individual Differences*, 25, 167-177.
- Seligman, M. E., e Csikszentmihalyi, M. (2000). Positive psychology. An introduction. *The American psychologist*, 55, 5-14.
- Seligman, M. E. (2002). Positive psychology, positive prevention, and positive therapy. *Handbook of positive psychology*, 2(2002), 3-12.
- Seligman, M. E., Steen, T. A., Park, N., e Peterson, C. (2005). Positive psychology progress: empirical validation of interventions. *The American psychologist*, 60, 410-421.
- Seligman, M. E., Ernst, R. M., Gillham, J., Reivich, K., e Linkins, M. (2009). Positive education: Positive psychology and classroom interventions. *Oxford review of education*, 35, 293-311.
- Steinberg L. (2005). Cognitive and affective development in adolescence. *Trends in cognitive sciences*, 9, 69-74.
- Tejada-Gallardo, C., Blasco-Belled, A., Torrelles-Nadal, C., e Alsinet, C. (2020). Effects of School-based Multicomponent Positive Psychology Interventions on Well-being and Distress in Adolescents: A Systematic Review and Meta-analysis. *Journal of youth and adolescence*, 49(10), 1943-1960.
- United Nations Children's Fund, The State of the World's Children 2021: *On My Mind – Promoting, protecting and caring for children's mental health*, UNICEF, New York, October 2021.
- Wagnild, G. M. (2009). *The Resilience Scale user's guide for the US English version of the Resilience Scale and the 14-item Resilience Scale (RS-14)*. The Resilience Center.
- Wood, A. M., Froh, J. J., e Geraghty, A. W. (2010). Gratitude and well-being: a review and theoretical integration. *Clinical psychology review*, 30, 890-905.
- World Health Organization. (1998). *Wellbeing measures in primary health care/the DepCare Project: report on a WHO meeting: Stockholm, Sweden, 12-13 February 1998* (No. WHO/EURO: 1998-4234-43993-62027). World Health Organization. Regional Office for Europe.